

ISIN

Reference List

Adams DC, Emerson RG, Heyer EJ, McCormick PC, Carmel PW, Stein BM, Farcy JP, Gallo EJ. Monitoring of intraoperative motor-evoked potentials under conditions of controlled neuromuscular blockade. *Anesth Analg* 1993;77(5):913-8.

Aglia LS, Romero R, Desai S, Ramirez M, Gonzalez AA, Gugino LD. The use of transcranial magnetic stimulation for monitoring descending spinal cord motor function. *Clin Electroencephalogr*. 2002;33(1):30-41.

Aglia LS, Kraus KH, Desai S, Romero R, Ramirez M, Levy WJ, Gugino LD. Efficacious use of a cap shaped coil for transcranial magnetic stimulation of descending motor paths. *Clin Electroencephalogr*. 2002;33(1):21-9.

Agnew WF, McCreery DB. Considerations for safety in the use of extracranial stimulation for motor evoked potentials. *Neurosurgery*. 1987;20(1):143-7.

Agnew WF, McCreery DB, Yuen TG, Bullara LA. Histologic and physiologic evaluation of electrically stimulated peripheral nerve: considerations for the selection of parameters. *Ann Biomed Eng*. 1989;17(1):39-60.

Agnew WF, McCreery DB, Yuen TG, Bullara LA. Evolution and resolution of stimulation-induced axonal injury in peripheral nerve. *Muscle Nerve*. 1999;22(10):1393-402.

Akagami R, Dong CC, Westerberg BD. Localized transcranial electrical motor evoked potentials for monitoring cranial nerves in cranial base surgery. *Neurosurgery*. 2005;57(1 Suppl):78-85.

Albright AL. Intraoperative spinal cord monitoring for intramedullary surgery: an essential adjunct? *Pediatr Neurosurg* 1998;29(2):112.

American Academy of Neurology. Assessment: intraoperative neurophysiology. Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology* 1990;40(11):1644-6.

American Electroencephalographic Society. Guideline eleven: guidelines for intraoperative monitoring of sensory evoked potentials. *J Clin Neurophysiol* 1994;11(1):77-87.

Aminoff MJ. Intraoperative monitoring by evoked potentials for spinal cord surgery: the cons. *Electroencephalogr Clin Neurophysiol* 1989;73(5):378-8

Andersson G, Ohlin A. Spatial facilitation of motor evoked responses in monitoring during spinal surgery. *Clin Neurophysiol* 1999;110(4):720-4.

Apel DM, Marrero G, King J, Tolo VT, Bassett GS. Avoiding paraplegia during anterior spinal surgery. The role of somatosensory evoked potential monitoring with temporary occlusion of segmental spinal arteries. *Spine* 1991;16(8 Suppl):S365-70.

Arrington ED, Hochschild DP, Steinagle TJ, Mongan PD, Martin SL. Monitoring of somatosensory and motor evoked potentials during open reduction and internal fixation of pelvis and acetabular fractures. *Orthopedics* 2000;23(10):1081-3.

Ashley RA, Wee AS. Intraoperative detection of unilateral spinal cord dysfunction by somatosensory evoked potentials. *J Miss State Med Assoc* 1988;29(11):339-41

Axon PR, Ramsden RT. Intraoperative electromyography for predicting facial function in vestibular schwannoma surgery. *Laryngoscope* 1999;109(6):922-6.

Bailey AR, Jones JG. Patients' memories of events during general anaesthesia. *Anaesthesia* 1997;52(5):460-76.

Balzer JR, Rose RD, Welch WC, Sclabassi RJ. Simultaneous somatosensory evoked potential and electromyographic recordings during lumbosacral decompression and instrumentation. *Neurosurgery* 1998;42(6):1318-24; discussion 1324-5.

Bar-On Z, Zeilig G, Blumen N, Ohry A, Azaria M. Paraplegia following surgical correction of scoliosis with Cotrel-Dubousset instrumentation. *Bull Hosp Jt Dis* 1995;54(1):32-4.

Bartley K, Woodforth IJ, Stephen JP, Burke D. Corticospinal volleys and compound muscle action potentials produced by repetitive transcranial stimulation during spinal surgery. *Clin Neurophysiol.* 2002;113(1):78-90.

Been HD, Kalkman CJ, Traast HS, Ongerboer de Visser BW. Neurologic injury after insertion of laminar hooks during Cotrel-Dubousset instrumentation. *Spine* 1994;19(12):1402-5.

Bejjani GK, Nora PC, Vera PL, Broemling L, Sekhar LN. The predictive value of intraoperative somatosensory evoked potential monitoring: review of 244 procedures. *Neurosurgery.* 1998;43(3):491-8; discussion 498-500.

Ben-David B, Haller G, Taylor P. Anterior spinal fusion complicated by paraplegia. A case report of a false-negative somatosensory-evoked potential. *Spine* 1987;12(6):536-9

Berardelli A, Inghilleri M, Cruccu G, Manfredi M. Corticospinal potentials after electrical and magnetic stimulation in man. *Electroencephalogr Clin Neurophysiol Suppl.* 1991;43:147-54.

Bernard JM, Pereon Y, Fayet G, Guiheneuc P. Effects of isoflurane and desflurane on neurogenic motor- and somatosensory-evoked potential monitoring for scoliosis surgery. *Anesthesiology* 1996;85(5):1013-9.

Binder DK, Lyon R, Manley GT. Transcranial motor evoked potential recording in a case of Kernohan's notch syndrome: case report. *Neurosurgery.* 2004;54(4):999-1002.

Boisseau N, Madany M, Staccini P, Armando G, Martin F, Grimaud D, Raucoules-Aime M. Comparison of the effects of sevoflurane and propofol on cortical somatosensory evoked potentials. *Br J Anaesth.* 2002;88(6):785-9.

Booke JS, Banta JV, Bunke FJ, Pelletier C. Somatosensory evoked potential monitoring during Cotrel-Dubousset instrumentation. Report of a case. *Spine* 1993;18(4):518-21.

Borrissov B, Langeron O, Lille F, Gomola A, Saillant G, Riou B, Viars P. Combination of propofol-sufentanil on somatosensory evoked potentials in surgery of the spine. *Ann Fr Anesth Reanim* 1995;14(4):326-30.

Bose B, Wierzbowski LR, Sestokas AK. Neurophysiologic monitoring of spinal nerve root function during instrumented posterior lumbar spine surgery. *Spine.* 2002;27(13):1444-50.

Bose B, Sestokas AK, Schwartz DM. Neurophysiological monitoring of spinal cord function during instrumented anterior cervical fusion. *Spine J.* 2004;4(2):202-7.

Bouchard JA, Bohlman HH, Biro C. Intraoperative improvements of somatosensory evoked potentials: correlation to clinical outcome in surgery for cervical spondylitic myelopathy. *Spine* 1996;21(5):589-94.

Boyd SG, Rothwell JC, Cowan JM, Webb PJ, Morley T, Asselman P, Marsden CD. A method of monitoring function in corticospinal pathways during scoliosis surgery with a note on motor conduction velocities. *J Neurol Neurosurg Psychiatry* 1986;49(3):251-7.

Bradford DS, Tay BK, Hu SS. Adult scoliosis: surgical indications, operative management, complications, and outcomes. *Spine* 1999;24(24):2617-29.

Bradshaw K, Webb JK, Fraser AM. Clinical evaluation of spinal cord monitoring in scoliosis surgery. *Spine* 1984;9(6):636-43.

Bridwell KH, Lenke LG, Baldus C, Blanke K. Major intraoperative neurologic deficits in pediatric and adult spinal deformity patients. Incidence and etiology at one institution. *Spine* 1998;23(3):324-31.

Brocke J, Irlbacher K, Hauptmann B, Voss M, Brandt SA. Transcranial magnetic and electrical stimulation compared: does TES activate intracortical neuronal circuits? *Clin Neurophysiol.* 2005;116(12):2748-56.

Brown RH, Nash CL Jr, Berilla JA, Amaddio MD. Cortical evoked potential monitoring. A system for intraoperative monitoring of spinal cord function. *Spine* 1984;9(3):256-61.

Burke D, Hicks RG, Stephen JP. Corticospinal volleys evoked by anodal and cathodal stimulation of the human motor cortex. *J Physiol.* 1990;425:283-99.

Burke D, Hicks R, Stephen J, Woodforth I, Crawford M. Assessment of corticospinal and somatosensory conduction simultaneously during scoliosis surgery. *Electroencephalogr Clin Neurophysiol* 1992;85(6):388-96.

Burke D, Hicks R, Stephen J, Woodforth I, Crawford M. Trial-to-trial variability of corticospinal volleys in human subjects. *Electroencephalogr Clin Neurophysiol*. 1995;97(5):231-7.

Burke D, Hicks RG. Surgical monitoring of motor pathways. *J Clin Neurophysiol* 1998;15(3):194-205.

Burke D, Nuwer MR, Daube J, Fischer C, Schramm J, Yingling CD, Jones SJ. Intraoperative monitoring. The International Federation of Clinical Neurophysiology. *Electroencephalogr Clin Neurophysiol Suppl* 1999;52:133-48.

Burke D, Bartley K, Woodforth IJ, Yakoubi A, Stephen JP. The effects of a volatile anaesthetic on the excitability of human corticospinal axons. *Brain* 2000;123 (Pt 5):992-1000.

Calancie B, Lebowitz N, Madsen P, Klose KJ. Intraoperative evoked EMG monitoring in an animal model. A new technique for evaluating pedicle screw placement. *Spine* 1992;17(10):1229-35.

Calancie B, Madsen P, Lebowitz N. Stimulus-evoked EMG monitoring during transpedicular lumbosacral spine instrumentation. Initial clinical results. *Spine* 1994;19(24):2780-6.

Calancie B, Harris W, Broton JG, Alexeeva N, Green BA. "Threshold-level" multipulse transcranial electrical stimulation of motor cortex for intraoperative monitoring of spinal motor tracts: description of method and comparison to somatosensory evoked potential monitoring. *J Neurosurg* 1998;88(3):457-70.

Calancie B, Harris W, Brindle GF, Green BA, Landy HJ. Threshold-level repetitive transcranial electrical stimulation for intraoperative monitoring of central motor conduction. *J Neurosurg (Spine 1)* 2001;95:161-8.

Carlioz H, Ouaknine M. Neurologic complications of surgery of the spine in children. *Chirurgie* 1994;120(11):26-30.

Cascino GD, So EL, Sharbrough FW, Strelow D, Lagerlund TD, Milde LN, O'Brien PC. Alfentanil-induced epileptiform activity in patients with partial epilepsy. *J Clin Neurophysiol*. 1993;10(4):520-5.

Cedzich C, Taniguchi M, Schafer S, Schramm J. Somatosensory evoked potential phase reversal and direct motor cortex stimulation during surgery in and around the central region. *Neurosurgery* 1996;38(5):962-70.

Cedzich C, Pechstein U, Schramm J, Schafer S. Electrophysiological considerations regarding electrical stimulation of motor cortex and brain stem in humans. *Neurosurgery*. 1998;42(3):527-32.

Cervellati S, Bettini N, Bianco T, Parisini P. Neurological complications in segmental spinal instrumentation: analysis of 750 patients. *Eur Spine J* 1996;5(3):161-6.

Chen Z. The effects of isoflurane and propofol on intraoperative neurophysiological monitoring during spinal surgery. *J Clin Monit Comput*. 2004;18(4):303-8.

Cheung AT, Pochettino A, McGarvey ML, Appoo JJ, Fairman RM, Carpenter JP, Moser WG, Woo EY, Bavaria JE. Strategies to manage paraplegia risk after endovascular stent repair of descending thoracic aortic aneurysms. *Ann Thorac Surg*. 2005;80(4):1280-8; discussion 1288-9.

Cioni B, Meglio M, Moles A, Tirendi M, Visocchi M. Spinal somatosensory evoked potential monitoring during microsurgery for syringomyelia: case reports. *Stereotact Funct Neurosurg* 1991;57(3):123-9.

Cioni B, Meglio M, Rossi GF. Intraoperative motor evoked potentials monitoring in spinal neurosurgery. *Arch Ital Biol* 1999;137(2-3):115-26.

Clapcich AJ, Emerson RG, Roye DP Jr, Xie H, Gallo EJ, Dowling KC, Ramnath B, Heyer EJ. The effects of propofol, small-dose isoflurane, and nitrous oxide on cortical somatosensory evoked potential and bispectral index monitoring in adolescents undergoing spinal fusion. *Anesth Analg*. 2004;99(5):1334-40.

Coles JG, Wilson GJ, Sima AF, Klement P, Tait GA. Intraoperative detection of spinal cord ischemia using somatosensory cortical evoked potentials during thoracic aortic occlusion. *Ann Thorac Surg* 1982;34(3):299-306.

Connolly JE. Hume Memorial lecture. Prevention of spinal cord complications in aortic surgery. *Am J Surg* 1998;176(2):92-101.

Crawford ES, Mizrahi EM, Hess KR, Coselli JS, Safi HJ, Patel VM. The impact of distal aortic perfusion and somatosensory evoked potential monitoring on prevention of paraplegia after aortic aneurysm operation. *J Thorac Cardiovasc Surg* 1988;95(3):357-67.

Crawford ES, Svensson LG, Hess KR, Shenaq SS, Coselli JS, Safi HJ, Mohindra PK, Rivera V. A prospective randomized study of cerebrospinal fluid drainage to prevent paraplegia after high-risk surgery on the thoracoabdominal aorta. *J Vasc Surg* 1991;13(1):36-45; discussion 45-6.

Cunningham JN Jr, Laschinger JC, Merkin HA, Nathan IM, Colvin S, Ransohoff J, Spencer FC. Measurement of spinal cord ischemia during operations upon the thoracic aorta: initial clinical experience. *Ann Surg*. 1982;196(3):285-96.

Cunningham JN Jr, Laschinger JC, Spencer FC. Monitoring of somatosensory evoked potentials during surgical procedures on the thoracoabdominal aorta. IV. Clinical observations and results. *J Thorac Cardiovasc Surg* 1987;94(2):275-85.

Darden BV 2nd, Hatley MK, Owen JH. Neurogenic motor evoked-potential monitoring in anterior cervical surgery. *J Spinal Disord* 1996;9(6):485-93.

Dasmahapatra HK, Coles JG, Taylor MJ, Cass D, Couper G, Adler S, Burrows F, Sherret H, Trusler GA, Williams WG. Identification of risk factors for spinal cord ischemia by the use of monitoring of somatosensory evoked potentials during coarctation repair. *Circulation*. 1987;76(3 Pt 2):III14-8.

Daube JR. Intraoperative monitoring by evoked potentials for spinal cord surgery: the pros. *Electroencephalogr Clin Neurophysiol* 1989;73(5):374-7.

Daube JR, Harper CM, Litchy WJ, Sharbrough FW. Intraoperative Monitoring. In: Daly DD and Pedley TA, eds. *Current practice of clinical electroencephalography*. 2nd ed. New York: Raven Press, 1990;739-79.

Dawson EG, Sherman JE, Kanim LE, Nuwer MR. Spinal cord monitoring. Results of the Scoliosis Research Society and the European Spinal Deformity Society survey. *Spine* 1991;16(8 Suppl):S361-4.

de Haan P, Kalkman CJ, Ubags LH, Jacobs MJ, Drummond JC. A comparison of the sensitivity of epidural and myogenic transcranial motor-evoked responses in the detection of acute spinal cord ischemia in the rabbit. *Anesth Analg* 1996;83(5):1022-7.

de Haan P, Kalkman CJ, de Mol BA, Ubags LH, Veldman DJ, Jacobs MJ. Efficacy of transcranial motor-evoked myogenic potentials to detect spinal cord ischemia during operations for thoracoabdominal aneurysms. *J Thorac Cardiovasc Surg*. 1997;113(1):87-100; discussion 100-1.

de Haan P, Kalkman CJ, Jacobs MJ. Spinal cord monitoring with myogenic motor evoked potentials: early detection of spinal cord ischemia as an integral part of spinal cord protective strategies during thoracoabdominal aneurysm surgery. *Semin Thorac Cardiovasc Surg* 1998;10(1):19-24.

de Haan P, Kalkman CJ, Meylaerts SA, Lips J, Jacobs MJ. Development of spinal cord ischemia after clamping of noncritical segmental arteries in the pig. *Ann Thorac Surg*. 1999;68(4):1278-84.

de Haan P, Kalkman CJ. Spinal cord monitoring: somatosensory- and motor-evoked potentials. *Anesthesiol Clin North America*. 2001;19(4):923-45.

de Mol B, Hamerlijnck R, Boezeman E, Vermeulen FE. Prevention of spinal cord ischemia in surgery of thoraco-abdominal aneurysms. The Bio Medicus pump, the recording of somatosensory evoked potentials and the impact on surgical strategy. *Eur J Cardiothorac Surg* 1990;4(12):658-64.

de Mol BA, Boezeman EH, Hamerlijnck RP, de Geest R. Experimental and clinical use of somatosensory evoked potentials in surgery of aneurysms of the descending thoracic aorta. *Thorac Cardiovasc Surg*. 1990;38(3):146-50.

de Noordhout AM, Born JD, Hans P, Remacle JM, Delwaide PJ. Intraoperative localisation of the primary motor cortex using single electrical stimuli. *J Neurol Neurosurg Psychiatry* 1996;60:442-444.

Delank KS, Delank HW, König DP, Popken F, Fürderer S, Eysel P. Iatrogenic paraplegia in spinal surgery. *Arch Orthop Trauma Surg.* 2005;125(1):33-41.

Delecrin J, Nguyen The Tich S, Passuti N, Pereaon Y. Neurogenic mixed evoked potential monitoring during scoliosis surgery: retrospective analysis of 149 cases. *Rev Chir Orthop Reparatrice Appar Mot* 2000;86(1):46-53.

Deletis V. Intraoperative Monitoring of the Functional Integrity of the Motor Pathways. In: Devinsky O, Beric A and Dogali M, Eds. *Electrical and Magnetic Stimulation of the Brain and Spinal Cord.* New York: Raven Press Ltd., 1993;201-14.

Deletis V, Kiproviski K, Morota N. The influence of halothane, enflurane, and isoflurane on motor evoked potentials. *Neurosurgery* 1993;33(1):173-4.

Deletis V, Morota N, Abbott IR. Electrodiagnosis in the management of brachial plexus surgery. *Hand Clin* 1995;11(4):555-61.

Deletis V, Rodi Z, Amassian VE. Neurophysiological mechanisms underlying motor evoked potentials in anesthetized humans. Part 2. Relationship between epidurally and muscle recorded MEPs in man. *Clin Neurophysiol* 2001;112(3):445-52.

Deletis V, Isgum V, Amassian VE. Neurophysiological mechanisms underlying motor evoked potentials in anesthetized humans. Part 1. Recovery time of corticospinal tract direct waves elicited by pairs of transcranial electrical stimuli. *Clin Neurophysiol* 2001;112(3):438-44.

Deletis V, Sala F. The role of intraoperative neurophysiology in the protection or documentation of surgically induced injury to the spinal cord. *Ann N Y Acad Sci* 2001;939:137-44.

Deletis V, Bueno De Camargo A. Interventional neurophysiological mapping during spinal cord procedures. *Stereotact Funct Neurosurg.* 2001;77(1-4):25-8.

Deletis V. Intraoperative neurophysiology and methodologies used to monitor the functional integrity of the motor system. In: Deletis V, Shils JL. eds. *Neurophysiology in Neurosurgery.* Academic Press, California. 2002;25-51.

Deletis V, Sala F. Intraoperative neurophysiological monitoring during spine surgery: an update. *Current Opinion in Orthopaedics* 2004;15:154–158.

DiCindio S, Theroux M, Shah S, Miller F, Dabney K, Brislin RP, Schwartz D. Multimodality monitoring of transcranial electric motor and somatosensory-evoked potentials during surgical correction of spinal deformity in patients with cerebral palsy and other neuromuscular disorders. *Spine* 2003;28(16):1851–1856.

Dinner DS, Luders H, Lesser RP, Morris HH, Barnett G, Klem G. Intraoperative spinal somatosensory evoked potential monitoring. *J Neurosurg* 1986;65(6):807-14.

Dobbs MB, Weinstein SL. Infantile and juvenile scoliosis. *Orthop Clin North Am* 1999;30(3):331-41.

Doita M, Marui T, Nishida K, Kurosaka M, Yoshiya S, Sha N. Anterior spinal artery syndrome after total spondylectomy of T10, T11, and T12. *Clin Orthop Relat Res*. 2002;(405):175-81.

Dolin MG. Late complications due to wire breakage in segmental spinal instrumentation. Report of two cases. *J Bone Joint Surg Am* 1984;66(6):967-8.

Dong C, MacDonald DB, Akagami R, Westerberg B, AlKhani A, AlShail E, Kanaan I, Hassounah M. Intraoperative facial motor evoked potential monitoring with transcranial electrical stimulation during skull base surgery. *Clin Neurophysiol* 2005;116:588-96.

Dong CC, MacDonald DB, Janusz MT. Intraoperative spinal cord monitoring during descending thoracic and thoracoabdominal aneurysm surgery. *Ann Thorac Surg*. 2002;74(5):S1873-6; discussion S1892-8.

Doty JR, Mahla ME, Furlow TW Jr. Intraoperative monitoring of evoked potentials with a spiral scalp electrode. Technical note. *Surg Neurol* 1988;30(1):73-4.

Draughn LR, Swanson TH. Axillary somatosensory evoked potential response: an alternate peripheral recording site. *J Clin Neurophysiol* 1998;15(1):64-8.

Drenger B, Parker SD, McPherson RW, North RB, Williams GM, Reitz BA, Beattie C. Spinal cord stimulation evoked potentials during thoracoabdominal aortic aneurysm surgery. *Anesthesiology* 1992;76(5):689-95.

Dudra J, Shiiya N, Matsui Y, Sakuma M, Ishii K, Asada M, Yasuda K. Operative results of thoracoabdominal repair for chronic type B aortic dissection. *J Cardiovasc Surg (Torino)* 1997;38(2):147-51.

Dunne JW, Field CM. The value of non-invasive spinal cord monitoring during spinal surgery and interventional angiography. *Clin Exp Neurol* 1991;28:199-209.

Dunne JW, Carroll WM. Intraoperative spinal cord monitoring. *Med J Aust* 1991;154(2):78-9.

Dunne JW, Silbert PL, Wren M. A prospective study of acute radiculopathy after scoliosis surgery. *Clin Exp Neurol* 1991;28:180-90.

Ecker ML, Dormans JP, Schwartz DM, Drummond DS, Bulman WA. Efficacy of spinal cord monitoring in scoliosis surgery in patients with cerebral palsy. *J Spinal Disord* 1996;9(2):159-64.

Elmore JR, Gloviczki P, Harper CM, Pairolero PC, Murray MJ, Bouchier RG, Bower TC, Daube JR. Failure of motor evoked potentials to predict neurologic outcome in experimental thoracic aortic occlusion. *J Vasc Surg*. 1991;14(2):131-9.

Elmore JR, Gloviczki P, Harper CM Jr, Murray MJ, Wu QH, Bower TC, Pairolero PC, Naessens JM, Daube JR. Spinal cord injury in experimental thoracic aortic occlusion: investigation of combined methods of protection. *J Vasc Surg*. 1992;15(5):789-98; discussion 798-9.

Erb TO, Ryhult SE, Duitmann E, Hasler C, Luetschg J, Frei FJ. Improvement of Motor-Evoked Potentials by Ketamine and Spatial Facilitation During Spinal Surgery in a Young Child. *Anesth Analg* 2005;100:1634–6.

Erwin CW, Erwin AC. Up and down the spinal cord: intraoperative monitoring of sensory and motor spinal cord pathways. *J Clin Neurophysiol* 1993;10(4):425-36.

Estrem SA, Haghghi S, Levy WJ, Wertheimer R, Kendall M. Motor-evoked potentials of facial musculature in dogs. *Laryngoscope* 1988;98(9):1012-5.

Ewen A, Bart BB, Goresky GV, Hoppenfeld S, Gross A, Lonner B. Correspondence. *The Journal of Bone and Joint Surgery* 1999;81:1044.

Faberowski LW, Black S, Trankina MF, Pollard RJ, Clark RK, Mahla ME. Somatosensory-evoked potentials during aortic coarctation repair. *J Cardiothorac Vasc Anesth*. 1999;13(5):538-43.

Fan D, Schwartz DM, Vaccaro AR, Hilibrand AS, Albert TJ. Intraoperative neurophysiologic detection of iatrogenic C5 nerve root injury during laminectomy for cervical compression myelopathy. *Spine*. 2002;27(22):2499-502.

Fava E, Bortolani EM, Ducati A, Ruberti U. Evaluation of spinal cord function by means of lower limb somatosensory evoked potentials in reparative aortic surgery. *J Cardiovasc Surg (Torino)*. 1988;29(4):421-7.

Firsching R, Heinen-Lauten M, Loeschke G. The effects of halothane and nitrous oxide on transcranial magnetic evoked potentials *Anesthesiol Intensivmed Notfallmed Schmerzther* 1991;26(7):381-3.

Fishback AS, Shields CB, Linden RD, Zhang YP, Burke D. The effects of propofol on rat transcranial magnetic motor evoked potentials. *Neurosurgery* 1995;37(5):969-74.

Fletcher JE, Hinn AR, Heard CM, Georges LS, Freid EB, Keifer A, Brooks SD, Bailey AG, Valley RD. The effects of isoflurane and desflurane titrated to a bispectral index of 60 on the cortical somatosensory evoked potential during pediatric scoliosis surgery. *Anesth Analg*. 2005;100(6):1797-803.

Floman Y, Micheli LJ, Penny JN, Riseborough EJ, Hall JE. Combined anterior and posterior fusion in seventy-three spinally deformed patients: indications, results and complications. *Clin Orthop* 1982;(164):110-22.

Florence G, Guerit JM, Gueguen B. Electroencephalography (EEG) and somatosensory evoked potentials (SEP) to prevent cerebral ischaemia in the operating room. *Neurophysiol Clin*. 2004;34(1):17-32.

Forbes HJ, Allen PW, Waller CS, Jones SJ, Edgar MA, Webb PJ, Ransford AO. Spinal cord monitoring in scoliosis surgery. Experience with 1168 cases. *J Bone Joint Surg Br* 1991;73(3):487-91.

Franco-Carcedo C, Villalibre I, Gonzalez-Hidalgo M. Protocol of intraoperative neurophysiological spinal cord monitoring. *Rev Neurol* 1999;28(6):583-7.

Fujiki M, Furukawa Y, Kamida T, Anan M, Inoue R, Abe T, Kobayashi H. Intraoperative corticomuscular motor evoked potentials for evaluation of motor function: a comparison with corticospinal D and I waves. *J Neurosurg*. 2006;104:85–92.

Fujioka H, Shimoji K, Tomita M, Denda S, Takada T, Homma T, Uchiyama S, Takahashi H, Tobita T, Baba H. Spinal cord potential recordings from the extradural space during scoliosis surgery. *Br J Anaesth* 1994;73(3):350-6.

Galla JD, Ergin MA, Sadeghi AM, Lansman SL, Danto J, Griep RB. A new technique using somatosensory evoked potential guidance during descending and thoracoabdominal aortic repairs. *J Card Surg* 1994;9(6):662-72.

Galla JD, Ergin MA, Lansman SL, McCullough JN, Nguyen KH, Spielvogel D, Klein JJ, Griep RB. Use of somatosensory evoked potentials for thoracic and thoracoabdominal aortic resections. *Ann Thorac Surg* 1999;67(6):1947-52; discussion 1953-8.

Galloway AC, Schwartz DS, Culliford AT, Ribakove GH, Grossi EA, Esposito RA, Baumann FG, Delianides J, Spencer FC, Colvin SB. Selective approach to descending thoracic aortic aneurysm repair: a ten-year experience. *Ann Thorac Surg* 1996;62(4):1152-7.

Gentili F, Lougheed WM, Yamashiro K, Corrado C. Monitoring of sensory evoked potentials during surgery of skull base tumours. *Can J Neurol Sci* 1985;12(4):336-40.

Ghaly RF, Lee JJ, Ham JH, Stone JL, George S, Raccforte P. Etomidate dose-response on somatosensory and transcranial magnetic induced spinal motor evoked potentials in primates. *Neurol Res* 1999;21(8):714-20.

Gharagozloo F, Neville RF Jr, Cox JL. Spinal cord protection during surgical procedures on the descending thoracic and thoracoabdominal aorta: a critical overview. *Semin Thorac Cardiovasc Surg* 1998;10(1):73-86.

Ghariani S, Spaey J, Liard L, Verhelst R, el Khoury G, Noirhomme P, d'Udekem Y, Matta A, Dion R, Guerit JM. Sensitivity, specificity, and impact on the surgical strategy of the perioperative neuromonitoring of somatic evoked potentials in vascular surgery performed with circulatory arrest under deep hypothermia. *Neurophysiol Clin* 1998;28(4):335-41.

Ghariani S, Liard L, Spaey J, Noirhomme PH, El Khoury GA, de Tourtchaninoff M, Dion RA, Guerit JM. Retrospective study of somatosensory evoked potential monitoring in deep hypothermic circulatory arrest. *Ann Thorac Surg* 1999;67(6):1915-8; discussion 1919-21.

Ghariani S, Matta A, Dion R, Guerit JM. Intra- and postoperative factors determining neurological complications after surgery under deep hypothermic

circulatory arrest: a retrospective somatosensory evoked potential study. *Clin Neurophysiol* 2000;111(6):1082-94.

Glassman SD, Johnson JR, Shields CB, Backman MH, Paloheimo MP, Edmonds HL Jr, Linden RD. Correlation of motor-evoked potentials, somatosensory-evoked potentials, and the wake-up test in a case of kyphoscoliosis. *J Spinal Disord* 1993;6(3):194-8.

Glassman SD, Shields CB, Linden RD, Zhang YP, Nixon AR, Johnson JR. Anesthetic effects on motor evoked potentials in dogs. *Spine* 1993;18(8):1083-9.

Glassman SD, Zhang YP, Shields CB, Johnson JR, Linden RD. Transcranial magnetic motor-evoked potentials in scoliosis surgery. *Orthopedics* 1995;18(10):1017-23.

Gloviczki P. Surgical repair of thoracoabdominal aneurysms: patient selection, techniques and results. *Cardiovasc Surg* 2002;10(4):434-441.

Gokaslan ZL, Samudrala S, Deletis V, Wildrick DM, Cooper PR. Intraoperative monitoring of spinal cord function using motor evoked potentials via transcutaneous epidural electrode during anterior cervical spinal surgery. *J Spinal Disord* 1997;10(4):299-303.

Goldbrunner RH, Schlake HP, Milewski C, Tonn JC, Helms J, Roosen K. Quantitative parameters of intraoperative electromyography predict facial nerve outcomes for vestibular schwannoma surgery. *Neurosurgery* 2000;46(5):1140-6; discussion 1146-8.

Gonzalez-Fajardo JA, Toledano M, Alvarez T, Vaquero C. Monitoring of evoked potentials during spinal cord ischaemia: experimental evaluation in a rabbit model. *Eur J Vasc Endovasc Surg*. 1998;16(4):320-8.

Grabitz K, Sandmann W, Stuhmeier K, Mainzer B, Godehardt E, Ohle B, Hartwich U. The risk of ischemic spinal cord injury in patients undergoing graft replacement for thoracoabdominal aortic aneurysms. *J Vasc Surg* 1996;23(2):230-40.

Griep RB, Ergin MA, Galla JD, Lansman S, Khan N, Quintana C, McCollough J, Bodian C. Looking for the artery of Adamkiewicz: a quest to minimize paraplegia after operations for aneurysms of the descending thoracic and thoracoabdominal aorta. *J Thorac Cardiovasc Surg* 1996;112(5):1202-13; discussion 1213-5.

Griep RB, Ergin MA, Galla JD, Klein JJ, Spielvogel D, Griep EB. Minimizing spinal cord injury during repair of descending thoracic and thoracoabdominal aneurysms: the Mount Sinai approach. *Semin Thorac Cardiovasc Surg* 1998;10(1):25-8.

Grubbs PE Jr, Marini C, Toporoff B, Nathan I, Basu S, Acinapura AJ, Cunningham JN Jr. Somatosensory evoked potentials and spinal cord perfusion pressure are significant predictors of postoperative neurologic dysfunction. *Surgery*. 1988;104(2):216-23.

Guerit JM, Verhelst R, Rubay J, Khoury G, Matta A, Dion R. Multilevel somatosensory evoked potentials (SEPs) for spinal cord monitoring in descending thoracic and thoraco-abdominal aortic surgery. *Eur J Cardiothorac Surg* 1996;10(2):93-103; discussion 103-4.

Guerit JM, Witdoeck C, Rubay J, Matta A, Dion R. The usefulness of the spinal and subcortical components of the posterior tibial nerve SEPs for spinal cord monitoring during aortic coarctation repair. *Electroencephalogr Clin Neurophysiol* 1997;104(2):115-21.

Guerit JM, Witdoeck C, de Tourtchaninoff M, Ghariani S, Matta A, Dion R, Verhelst R. Somatosensory evoked potential monitoring in carotid surgery. I. Relationships between qualitative SEP alterations and intraoperative events. *Electroencephalogr Clin Neurophysiol* 1997;104(6):459-69.

Guerit JM. New methods on the subject of peroperative neurophysiological monitoring in vascular and orthopedic surgery. *Bull Mem Acad R Med Belg* 1998;153(7-9):325-31; discussion 332-3.

Guerit JM, Witdoeck C, Verhelst R, Matta AJ, Jacquet LM, Dion RA. Sensitivity, specificity, and surgical impact of somatosensory evoked potentials in descending aorta surgery. *Ann Thorac Surg* 1999;67(6):1943-6; discussion 1953-8.

Guerit JM, Dion RA. State-of-the-art of neuromonitoring for prevention of immediate and delayed paraplegia in thoracic and thoracoabdominal aorta surgery. *Ann Thorac Surg*. 2002;74(5):S1867-9; discussion S1892-8.

Gugino LD, Kraus KH, Heino R, Aglio LS, Levy WJ, Cohn L, Maddi R. Peripheral ischemia as a complicating factor during somatosensory and motor evoked potential monitoring of aortic surgery. *J Cardiothorac Vasc Anesth* 1992;6(6):715-9.

Gundanna M, Eskenazi M, Bendo J, Spivak J, Moskovich R. Somatosensory evoked potential monitoring of lumbar pedicle screw placement for in situ posterior spinal fusion. *Spine J*. 2003;3(5):370-6.

Gunnarsson T, Krassioukov AV, Sarjeant R, Fehlings MG. Real-time continuous intraoperative electromyographic and somatosensory evoked potential recordings in spinal surgery: correlation of clinical and electrophysiologic findings in a prospective, consecutive series of 213 cases. *Spine*. 2004;29(6):677-84.

Haghighi SS, York DH, Ebeling J, Gumerlock MK, Oro JJ, Gaines RW. Dissociation of somatosensory and motor evoked potentials in a patient with an intramedullary spinal tumor. *Mo Med* 1992;89(11):790-4.

Haghighi SS, York DH, Gaines RW, Oro JJ. Monitoring of motor tracts with spinal cord stimulation. *Spine* 1994;19(13):1518-24.

Hamaya H. Study on prevention of paraplegia during occlusion blockade of the thoracic aorta--examination of the evoked potential in the motor nerve induced by stimulation of the motor area of cerebral cortex. *Nippon Kyobu Geka Gakkai Zasshi*. 1993;41(8):1347-56.

- Hargreaves SJ, Watt JW. Intravenous anaesthesia and repetitive transcranial magnetic stimulation monitoring in spinal column surgery. *Br J Anaesth*. 2005;94(1):70-3.
- Harper CM Jr, Daube JR, Litchy WJ, Klassen RA. Lumbar radiculopathy after spinal fusion for scoliosis. *Muscle Nerve* 1988;11(4):386-91.
- Harper CM, Daube JR. Facial nerve electromyography and other cranial nerve monitoring. *J Clin Neurophysiol* 1998;15(3):206-16.
- Haupt WF, Erasmi-Korber H, Lanfermann H. Intraoperative recording of parietal SEP can miss hemodynamic infarction during carotid endarterectomy: a case study. *Electroencephalogr Clin Neurophysiol* 1994;92(1):86-8.
- Hausmann ON, Minb K, Boosb N, Ruetschc YA, Ernia T, Curta A. Transcranial electrical stimulation: significance of fast versus slow charge delivery for intraoperative monitoring. *Clin Neurophysiol*. 2002;113:1532–1535.
- Hays SR, Schwengel DA. Transient hypotension as a complication of monitoring transcervical motor evoked potentials. *Anesthesiology* 1999;90(1):314-7.
- Helmets SL, Hall JE. Intraoperative somatosensory evoked potential monitoring in pediatrics. *J Pediatr Orthop*. 1994;14(5):592-8.
- Herdmann J, Lumenta CB, Huse KO. Magnetic stimulation for monitoring of motor pathways in spinal procedures. *Spine* 1993;18(5):551-9.
- Herdmann J, Deletis V, Edmonds HL Jr, Morota N. Spinal cord and nerve root monitoring in spine surgery and related procedures. *Spine* 1996;21(7):879-85.
- Hicks R, Burke D, Stephen J, Woodforth I, Crawford M. Corticospinal volleys evoked by electrical stimulation of human motor cortex after withdrawal of volatile anaesthetics. *J Physiol*. 1992;456:393-404.
- Hicks RG, Burke DJ, Stephen JP. Monitoring spinal cord function during scoliosis surgery with Cotrel-Dubousset instrumentation. *Med J Australia* 1991;154:82-86.
- Hicks RG, Woodforth IJ, Crawford MR, Stephen JP, Burke DJ. Some effects of isoflurane on I waves of the motor evoked potential. *Br J Anaesth*. 1992;69(2):130-6.
- Hilibrand AS, Schwartz DM, Sethuraman V, Vaccaro AR, Albert TJ. Comparison of transcranial electric motor and somatosensory evoked potential monitoring during cervical spine surgery. *J Bone Joint Surg Am*. 2004;86-A(6):1248-53.
- Holdefer RN, Sadleir R, Russell MJ. Predicted current densities in the brain during transcranial electrical stimulation. *Clin Neurophysiol*. 2006;117(6):1388-97.
- Holland NR, Kostuik JP. Continuous electromyographic monitoring to detect nerve root injury during thoracolumbar scoliosis surgery. *Spine* 1997;22(21):2547-50.

Holland NR. Subcortical strokes from intracranial aneurysm surgery: implications for intraoperative neuromonitoring. *J Clin Neurophysiol.* 1998;15(5):439-46.

Hone SW, Commins DJ, Rames P, Chen JM, Rowed D, McLean A, Nedzelski JM. Prognostic factors in intraoperative facial nerve monitoring for acoustic neuroma. *J Otolaryngol* 1997;26(6):374-8.

Horikoshi T, Omata T, Uchida M, Asari Y, Nukui H. Usefulness and pitfalls of intraoperative spinal motor evoked potential recording by direct cortical electrical stimulation. *Acta Neurochir (Wien)* 2000;142(3):257-62.

Horiuchi K, Suzuki K, Sasaki T, Matsumoto M, Sakuma J, Konno Y, Oinuma M, Itakura T, Kodama N. Intraoperative monitoring of blood flow insufficiency during surgery of middle cerebral artery aneurysms. *J Neurosurg.* 2005;103(2):275-83.

Hurlbert RJ, Fehlings MG, Moncada MS. Use of sensory-evoked potentials recorded from the human occiput for intraoperative physiologic monitoring of the spinal cord. *Spine* 1995;20(21):2318-27.

Hurlbert RJ, Tator CH, Fehlings MG, Koyanagi I, Moncada MS. A novel sensory evoked potential for selective monitoring of the ventral spinal cord: from bench to bedside. *Electroencephalogr Clin Neurophysiol Suppl* 1996;46:221-31

Ihaya A, Morioka K, Noguchi H, Kimura T, Nishii H, Hiramatsu Y, Chiba Y, Muraoka R. A case report of descending thoracic aortic aneurysm associated with anterior spinal artery syndrome despite no marked ESP changes. *Kyobu Geka* 1990;43(10):843-6.

Inghilleri M, Berardelli A, Cruccu G, Priori A, Manfredi M. Corticospinal potentials after transcranial stimulation in humans. *J Neurol Neurosurg Psychiatry.* 1989;52(8):970-4.

Inoue S, Kawaguchi M, Takashi S, Kakimoto M, Sakamoto T, Kitaguchi K, Furuya H, Morimoto T, Sakaki T. Intraoperative monitoring of myogenic motor-evoked potentials from the external anal sphincter muscle to transcranial electrical stimulation. *Spine.* 2002;27(21):E454-9.

Iwasaki H, Tamaki T, Yoshida M, Ando M, Yamada H, Tsutsui S, Takami M. Efficacy and limitations of current methods of intraoperative spinal cord monitoring. *J Orthop Sci.* 2003;8(5):635-42.

Jacobs MJ, Meylaerts SA, de Haan P, de Mol BA, Kalkman CJ. Strategies to prevent neurologic deficit based on motor-evoked potentials in type I and II thoracoabdominal aortic aneurysm repair. *J Vasc Surg* 1999;29(1):48-57; discussion 57-9.

Jacobs MJ, Meylaerts SA, de Haan P, de Mol BA, Kalkman CJ. Assessment of spinal cord ischemia by means of evoked potential monitoring during thoracoabdominal aortic surgery. *Semin Vasc Surg* 2000;13(4):299-307.

Jacobs MJ, de Mol BA, Elenbaas T, Mess WH, Kalkman CJ, Schurink GW, Mochtar B. Spinal cord blood supply in patients with thoracoabdominal aortic aneurysms. *J Vasc Surg.* 2002;35(1):30-7.

Jacobs MJ, Elenbaas TW, Schurink GW, Mess WH, Mochtar B. Assessment of spinal cord integrity during thoracoabdominal aortic aneurysm repair. *Ann Thorac Surg.* 2002;74(5):S1864-6; discussion S1892-8.

Jacobs MJ, Mess WH. The role of evoked potential monitoring in operative management of type I and type II thoracoabdominal aortic aneurysms. *Semin Thorac Cardiovasc Surg* 2003;15(4):353-64.

Jellinek D, Platt M, Jewkes D, Symon L. Effects of nitrous oxide on motor evoked potentials recorded from skeletal muscle in patients under total anesthesia with intravenously administered propofol. *Neurosurgery* 1991;29(4):558-62.

Jellinek D, Jewkes D, Symon L. Noninvasive intraoperative monitoring of motor evoked potentials under propofol anesthesia: effects of spinal surgery on the amplitude and latency of motor evoked potentials. *Neurosurgery* 1991;29(4):551-7.

Jimenez JC, Sani S, Braverman B, Deutsch H, Ratliff JK. Palsies of the fifth cervical nerve root after cervical decompression: prevention using continuous intraoperative electromyography monitoring. *J Neurosurg Spine.* 2005;3(2):92-7.

John ER, Chabot RJ, Prichep LS, Ransohoff J, Epstein F, Berenstein A. Real-time intraoperative monitoring during neurosurgical and neuroradiological procedures. *J Clin Neurophysiol.* 1989;6(2):125-58.

Johnston CE 2nd, Happel LT Jr, Norris R, Burke SW, King AG, Roberts JM. Delayed paraplegia complicating sublaminar segmental spinal instrumentation. *J Bone Joint Surg Am* 1986;68(4):556-63.

Jones SC, Fernau R, Woeltjen BL. Use of somatosensory evoked potentials to detect peripheral ischemia and potential injury resulting from positioning of the surgical patient: case reports and discussion. *Spine J.* 2004;4(3):360-2.

Jones SJ, Edgar MA, Ransford AO, Thomas NP. A system for the electrophysiological monitoring of the spinal cord during operations for scoliosis. *J Bone Joint Surg Br* 1983;65(2):134-9.

Jones SJ, Harrison R, Koh KF, Mendoza N, Crockard HA. Motor evoked potential monitoring during spinal surgery: responses of distal limb muscles to transcranial cortical stimulation with pulse trains. *Electroencephalogr Clin Neurophysiol* 1996;100(5):375-83.

Jones SJ, Buonamassa S, Crockard HA. Two cases of quadriplegia following anterior cervical discectomy, with normal perioperative somatosensory evoked potentials. *J Neurol Neurosurg Psychiatry.* 2003;74(2):273-6.

Journee HL, Polak HE, de Kleuver M. Influence of electrode impedance on threshold voltage for transcranial electrical stimulation in motor evoked potential monitoring. *Med Biol Eng Comput.* 2004;42(4):557-61.

Kai Y, Owen JH, Lenke LG, Bridwell KH, Oakley DM, Sugioka Y. Use of sciatic neurogenic motor evoked potentials versus spinal potentials to predict early-onset neurologic deficits when intervention is still possible during overdistracted. *Spine* 1993;18(9):1134-9.

Kai Y, Owen JH, Allen BT, Dobras M, Davis C. Relationship between evoked potentials and clinical status in spinal cord ischemia. *Spine* 1994;19(10):1162-7.

Kakimoto M, Inoue S, Sakamoto T, Kawaguchi M, Kitaguchi K, Furuya H. Intraoperative motor evoked potential monitoring: a review of 115 cases. *Masui* 2000;49(3):240-4.

Kakinohana M, Kawabata T, Miyata Y, Sugahara K. Myogenic transcranial motor evoked potentials monitoring cannot always predict neurologic outcome after spinal cord ischemia in rats. *J Thorac Cardiovasc Surg* 2005;129:46-52.

Kalita J, Srivastava AK, Mittal P, Sharma VP, Misra UK. Evoked potential changes in ischaemic myelopathy. *Electromyogr Clin Neurophysiol.* 2003;43(4):211-5.

Kalkman CJ, ten Brink SA, Been HD, Bovill JG. Variability of somatosensory cortical evoked potentials during spinal surgery. Effects of anesthetic technique and high-pass digital filtering. *Spine* 1991;16(8):924-9.

Kalkman CJ, Drummond JC, Ribberink AA. Low concentrations of isoflurane abolish motor evoked responses to transcranial electrical stimulation during nitrous oxide/opioid anesthesia in humans. *Anesth Analg* 1991;73(4):410-5.

Kalkman CJ, Romijn K, Denslagen W. Eliminating diathermy-induced artifacts during intraoperative monitoring of somatosensory-evoked potentials: a hardware solution. *J Clin Monit* 1991;7(4):320-4.

Kalkman CJ, Traast H, Zuurmond WW, Bovill JG. Differential effects of propofol and nitrous oxide on posterior tibial nerve somatosensory cortical evoked potentials during alfentanil anaesthesia. *Br J Anaesth* 1991;66(4):483-9.

Kalkman CJ, Drummond JC, Ribberink AA, Patel PM, Sano T, Bickford RG. Effects of propofol, etomidate, midazolam, and fentanyl on motor evoked responses to transcranial electrical or magnetic stimulation in humans. *Anesthesiology* 1992;76(4):502-9.

Kalkman CJ, Drummond JC, Kennelly NA, Patel PM, Partridge BL. Intraoperative monitoring of tibialis anterior muscle motor evoked responses to transcranial electrical stimulation during partial neuromuscular blockade. *Anesth Analg* 1992;75(4):584-9.

Kalkman CJ, Been HD, Ongerboer de Visser BW. Intraoperative monitoring of spinal cord function. A review. *Acta Orthop Scand* 1993;64(1):114-23.

Kalkman CJ, Drummond JC, Patel PM, Sano T, Chesnut RM. Effects of droperidol, pentobarbital, and ketamine on myogenic transcranial magnetic motor-evoked responses in humans. *Neurosurgery* 1994;35(6):1066-71.

Kaneko K, Kawai S, Taguchi T, Fuchigami Y, Morita H, Ofuji A, Yonemura H. Spatial distribution of corticospinal potentials following transcranial electric and magnetic stimulation in human spinal cord. *J Neurol Sci.* 1997;151(2):217-21.

Kaneko M, Fukamachi A, Sasaki H, Miyazawa N, Yagishita T, Nukui H. Intraoperative monitoring of the motor function: experimental and clinical study. *Acta Neurochir Suppl (Wien)* 1988;42:18-21.

Kaplan BJ, Friedman WA, Alexander JA, Hampson SR. Somatosensory evoked potential monitoring of spinal cord ischemia during aortic operations. *Neurosurgery.* 1986;19(1):82-90.

Kaplan BJ, Friedman WA, Gravenstein N, Richards R, Davis RF. Effects of aortic occlusion on regional spinal cord blood flow and somatosensory evoked potentials in sheep. *Neurosurgery.* 1987;21(5):668-75.

Kartush JM. Intra-operative monitoring in acoustic neuroma surgery. *Neurol Res* 1998;20(7):593-6.

Kawaguchi M, Sakamoto T, Ohnishi H, Shimizu K, Karasawa J, Furuya H. Intraoperative myogenic motor evoked potentials induced by direct electrical stimulation of the exposed motor cortex under isoflurane and sevoflurane. *Anesth Analg* 1996;82(3):593-9.

Kawaguchi M, Sakamoto T, Inoue S, Kakimoto M, Furuya H, Morimoto T, Sakaki T. Low dose propofol as a supplement to ketamine-based anesthesia during intraoperative monitoring of motor-evoked potentials. *Spine* 2000;25(8):974-9.

Keller BP, Haghghi SS, Oro JJ, Eggers GW Jr. The effects of propofol anesthesia on transcortical electric evoked potentials in the rat. *Neurosurgery* 1992;30(4):557-60.

Kerz T, Hennes HJ, Fève A, Decq P, Filipetti P, Duvaldestin P. Effects of propofol on H-reflex in humans. *Anesthesiology.* 2001;94(1):32-7.

Kimovec MA, Koht A, Sloan TB. Effects of sufentanil on median nerve somatosensory evoked potentials. *Br J Anaesth* 1990;65(2):169-72.

Kitagawa H, Takano H, Takakuwa K, Yamamoto N, Kida Y, Tsuji H. Origins and conducting pathways of motor evoked potentials elicited by transcranial (vertex-hard palate) stimulation in cats. *Neurosurgery.* 1991;28(3):358-63.

Kitagawa H, Nakamura H, Kawaguchi Y, Tsuji H, Satone T, Takano H, Nakatoh S. Magnetic-evoked compound muscle action potential neuromonitoring in spine surgery. *Spine* 1995;20(20):2233-9.

Kobara N, Owen JH, Kostuik J, Huckell C, Tooke SM. Relative effectiveness of electrically- vs mechanically-elicited EMGs in detecting pedicle wall perforation and surgically-induced nerve root damage. *Fukuoka Igaku Zasshi* 2000;91(2):40-54.

Kochs E, Kalkman CJ, Thornton C, Newton D, Bischoff P, Kuppe H, Abke J, Konecny E, Nahm W, Stockmanns G. Middle latency auditory evoked responses and electroencephalographic derived variables do not predict movement to noxious stimulation during 1 minimum alveolar anesthetic concentration isoflurane/nitrous oxide anesthesia. *Anesth Analg* 1999;88(6):1412-7.

Koht A, Schutz W, Schmidt G, Schramm J, Watanabe E. Effects of etomidate, midazolam, and thiopental on median nerve somatosensory evoked potentials and the additive effects of fentanyl and nitrous oxide. *Anesth Analg* 1988;67(5):435-41.

Koizumi N, Obitsu Y, Koide K, Sato K, Saiki N, Watanabe Y, Ichihashi H, Yokoi Y, Shimazaki T, Kawaguchi S, Ishimaru S. Evaluation of spinal cord ischemia in endovascular stent graft repair and surgical operation of descending thoracic or thoracoabdominal aortic aneurysms. *Kyobu Geka*. 2004;57(4):262-7.

Kolenda H, Steffens H, Gefeller O, Hagenah J, Schomburg ED. Critical levels of spinal cord blood flow and duration of ischemia for the acute recovery of segmental spinal cord responses in cats. *J Spinal Disord*. 1997;10(4):288-95.

Komanetsky RM, Padberg AM, Lenke LG, Bridwell KH, Russo MH, Chapman MP, Hamill CL. Neurogenic motor evoked potentials: a prospective comparison of stimulation methods in spinal deformity surgery. *J Spinal Disord* 1998;11(1):21-8.

Kombos T, Suess O, Funk T, Kern BC, Brock M. Intra-operative mapping of the motor cortex during surgery in and around the motor cortex. *Acta Neurochir (Wien)* 2000;142(3):263-8.

Kombos T, Suess O, Pietila T, Brock M. Subdural air limits the elicitation of compound muscle action potentials by high-frequency transcranial electrical stimulation. *Br J Neurosurg* 2000;14(3):240-3.

Kombos T, Suess O, Ciklatekerlio O, Brock M. Monitoring of intraoperative motor evoked potentials to increase the safety of surgery in and around the motor cortex. *J Neurosurg*. 2001;95(4):608-14.

Kondo R, Saito S, Kuroki A, Sato S, Katakura K, Kayama T. Significance and usefulness of corticospinal motor evoked potential monitoring for lesions adjacent to primary motor cortex. *No To Shinkei*. 2004;56(6):496-502.

Kothbauer K, Schmid UD, Seiler RW, Eisner W. Intraoperative motor and sensory monitoring of the cauda equina. *Neurosurgery* 1994;34(4):702-7; discussion 707.

Kothbauer K, Deletis V, Epstein FJ. Intraoperative spinal cord monitoring for intramedullary surgery: an essential adjunct. *Pediatr Neurosurg* 1997;26(5):247-54.

Kothbauer KF, Deletis V, Epstein FJ. Intraoperative monitoring. *Pediatr Neurosurg* 1998;29(1):54-5.

Kothbauer KF. Motor evoked potential monitoring for intramedullary spinal cord tumor surgery. In: Deletis V, Shils JL. eds. Neurophysiology in Neurosurgery. Academic Press, California 2002;73-92.

Kothbauer KF, Novak K. Intraoperative monitoring for tethered cord surgery: an update. Neurosurg Focus. 2004;16(2):E8.

Koyanagi I, Iwasaki Y, Isu T, Abe H, Akino M, Kuroda S. Spinal cord evoked potential monitoring after spinal cord stimulation during surgery of spinal cord tumors. Neurosurgery 1993;33(3):451-9; discussion 459-60.

Kraus KH, Pope ER, O'Brien D, Hay BL. The effects of aortic occlusion on transcranially induced evoked potentials in the dog. Vet Surg. 1990;19(5):341-7.

Kraus KH, Gugino LD, Levy WJ, Cadwell J, Roth BJ. The use of a cap-shaped coil for transcranial magnetic stimulation of the motor cortex. J Clin Neurophysiol. 1993;10(3):353-62.

Krieger D, Adams HP, Albert F, von Haken M, Hacke W. Pure motor hemiparesis with stable somatosensory evoked potential monitoring during aneurysm surgery: case report. Neurosurgery 1992;31(1):145-50.

Krieger D, Sciabassi RJ. Time-varying evoked potentials. J Med Eng Technol. 1994;18(3):96-100.

Krodel A, Rehmet JC, Hamburger C. Spinal cord compression caused by the rod of a Harrington instrumentation device: a late complication in scoliosis surgery. Eur Spine J 1997;6(3):208-10.

Krombach GA, Spetzger U, Rohde V, Gilsbach JM. Intraoperative localization of functional regions in the sensorimotor cortex by neuronavigation and cortical mapping. Comput Aided Surg 1998;3(2):64-73.

Ku AS, Hu Y, Irwin MG, Chow B, Gunawardene S, Tan EE, Luk KD. Effect of sevoflurane/nitrous oxide versus propofol anaesthesia on somatosensory evoked potential monitoring of the spinal cord during surgery to correct scoliosis. Br J Anaesth. 2002;88(4):502-7.

Kumar A, Bhattacharya A, Makhija N. Evoked potential monitoring in anaesthesia and analgesia. Anaesthesia 2000;55(3):225-41.

Kunihara T, Shiiya N, Yasuda K. Strategy for spinal cord protection during thoracoabdominal aortic surgery. Kyobu Geka. 2004;57(4):319-24.

Labrom RD, Hoskins, Reilly CW, Tredwell SJ, Wong, PKH. Clinical Usefulness of Somatosensory Evoked Potentials for Detection of Brachial Plexopathy Secondary to Malpositioning in Scoliosis Surgery. Spine 30(18):2089-2093.

Lalwani AK, Butt FY, Jackler RK, Pitts LH, Yingling CD. Delayed onset facial nerve dysfunction following acoustic neuroma surgery. Am J Otol 1995;16(6):758-64.

LaMont RL, Wasson SL, Green MA. Spinal cord monitoring during spinal surgery using somatosensory spinal evoked potentials. *J Pediatr Orthop* 1983;3(1):31-6.

Lang EW, Beutler AS, Chesnut RM, Patel PM, Kennelly NA, Kalkman CJ, Drummond JC, Garfin SR. Myogenic motor-evoked potential monitoring using partial neuromuscular blockade in surgery of the spine. *Spine* 1996;21(14):1676-86.

Langeloo DD, Lelivelt A, Louis Journée H, Slappendel R, de Kleuver M. Transcranial electrical motor-evoked potential monitoring during surgery for spinal deformity: a study of 145 patients. *Spine*. 2003;28(10):1043-50.

Langeloo DD, Lelivelt A, Journee HL, Slappendel R, de Kleuver M. Transcranial electrical motor-evoked potential monitoring during surgery for spinal deformity: A study of 145 patients. *Spine* 2003;28(10):1043–1050.

Langeron O, Lille F, Zerhouni O, Orliaguet G, Saillant G, Riou B, Coriat P. Comparison of the effects of ketamine-midazolam with those of fentanyl-midazolam on cortical somatosensory evoked potentials during major spine surgery. *Br J Anaesth* 1997;78(6):701-6.

Langeron O, Vivien B, Paqueron X, Saillant G, Riou B, Coriat P, Lille F. Effects of propofol, propofol-nitrous oxide and midazolam on cortical somatosensory evoked potentials during sufentanil anaesthesia for major spinal surgery. *Br J Anaesth* 1999;82(3):340-5.

Laschinger JC, Cunningham JN Jr, Isom OW, Nathan IM, Spencer FC. Definition of the safe lower limits of aortic resection during surgical procedures on the thoracoabdominal aorta: use of somatosensory evoked potentials. *J Am Coll Cardiol* 1983;2(5):959-65.

Laschinger JC, Cunningham JN Jr, Baumann FG, Isom OW, Spencer FC. Monitoring of somatosensory evoked potentials during surgical procedures on the thoracoabdominal aorta. II. Use of somatosensory evoked potentials to assess adequacy of distal aortic bypass and perfusion after thoracic aortic cross-clamping. *J Thorac Cardiovasc Surg* 1987;94:266-270.

Laschinger JC, Cunningham JN Jr, Baumann FG, Cooper MM, Krieger KH, Spencer FC. Monitoring of somatosensory evoked potentials during surgical procedures on the thoracoabdominal aorta. III. Intraoperative identification of vessels critical to spinal cord blood supply. *J Thorac Cardiovasc Surg* 1987;94:271-274.

Laschinger JC, Owen J, Rosenbloom M, Cox JL, Kouchoukos NT. Direct noninvasive monitoring of spinal cord motor function during thoracic aortic occlusion: use of motor evoked potentials. *J Vasc Surg* 1988;7(1):161-71.

Laureau E, Marciniak B, Hebrard A, Herbaux B, Guieu JD. Neuromonitoring and anesthesia in surgery of the spine. *Neurophysiol Clin* 1998;28(4):299-320.

Laureau E, Marciniak B, Hebrard A, Herbaux B, Guieu JD. Comparative study of propofol and midazolam effects on somatosensory evoked potentials during surgical treatment of scoliosis. *Neurosurgery* 1999;45(1):69-74; discussion 75.

Lee WY, Hou WY, Yang LH, Lin SM. Intraoperative monitoring of motor function by magnetic motor evoked potentials. *Neurosurgery* 1995;36(3):493-500.

Legatt AD, Ellen R. Grass Lecture: Motor evoked potential monitoring. *Am J Electroneurodiagnostic Technol.* 2004;44(4):223-43.

Leis AA, Zhou HH, Mehta M, Harkey HL 3rd, Paske WC. Behavior of the H-reflex in humans following mechanical perturbation or injury to rostral spinal cord. *Muscle Nerve.* 1996;19(11):1373-82.

Leppanen R, Maguire J, Wallace S, Madigan R, Draper V. Intraoperative lower extremity reflex muscle activity as an adjunct to conventional somatosensory-evoked potentials and descending neurogenic monitoring in idiopathic scoliosis. *Spine* 1995;20(17):1872-7.

Leppanen R, Madigan R, Sears C, Maguire J, Wallace S, Captain J. Intraoperative collision studies demonstrate descending spinal cord stimulated evoked potentials and ascending somatosensory evoked potentials are mediated through common pathways (abstr.). *J Clin Neurophysiology* 1999;16:170.

Leppanen RE, American Society of Neurophysiological Monitoring. Intraoperative monitoring of segmental spinal nerve root function with free-run and electrically-triggered electromyography and spinal cord function with reflexes and F-responses. A position statement by the American Society of Neurophysiological Monitoring. *J Clin Monit Comput.* 2005;19(6):437-61. Epub 2006 Jan 25.

Lesser RP, Lueders H, Hahn J, Klem G. Early somatosensory potentials evoked by median nerve stimulation: intraoperative monitoring. *Neurology* 1981;31(12):1519-23.

Lesser RP, Raudzens P, Luders H, Nuwer MR, Goldie WD, Morris HH 3rd, Dinner DS, Klem G, Hahn JF, Shetter AG, et al. Postoperative neurological deficits may occur despite unchanged intraoperative somatosensory evoked potentials. *Ann Neurol* 1986;19(1):22-5.

Levy JW. Clinical experience with motor and cerebellar evoked potential monitoring. *Neurosurgery* 1987;20:169-182.

Levy WJ. Spinal evoked potentials from the motor tracts. *J Neurosurg* 1983;58(1):38-44.

Levy WJ, York DH, McCaffrey M, Tanzer F. Motor evoked potentials from transcranial stimulation of the motor cortex in humans. *Neurosurgery* 1984;15(3):287-302.

Levy WJ, McCaffrey M, York DH, Tanzer F. Motor evoked potentials from transcranial stimulation of the motor cortex in cats. *Neurosurgery.* 1984;15(2):214-27.

Levy WJ. Transcranial stimulation of the motor cortex to produce motor-evoked potentials. *Med Instrum* 1987;21(5):248-54.

Linstedt U, Maier C, Petry A. Intraoperative monitoring with somatosensory evoked potentials in carotid artery surgery--less reliable in patients with preoperative neurologic deficiency? *Acta Anaesthesiol Scand*. 1998;42(1):13-6.

Lips J, De Haan P, Bouma GJ, Jacobs MJ, Kalkman CJ. Delayed detection of motor pathway dysfunction after selective reduction of thoracic spinal cord blood flow in pigs. *J Thorac Cardiovasc Surg*. 2002;123(3):531-538.

Liscic RM, Morota N, Deletis V. Intramedullar stimulation of the facial and hypoglossal nerves: estimation of the stimulated site. *Croat Med J* 2000;41(4):384-8.

Lorenzini NA, Schneider JH. Temporary loss of intraoperative motor-evoked potential and permanent loss of somatosensory-evoked potentials associated with a postoperative sensory deficit. *J Neurosurg Anesthesiol* 1996;8(2):142-7.

Lubitz SE, Keith RW, Crawford AH. Intraoperative experience with neuromotor evoked potentials. A review of 60 consecutive cases. *Spine* 1999;24(19):2030-3.

Lueders H, Gurd A, Hahn J, Andrich J, Weiker G, Klem G. A new technique for intraoperative monitoring of spinal cord function: multichannel recording of spinal cord and subcortical evoked potentials. *Spine* 1982;7(2):110-5.

Luk KD, Hu Y, Wong YW, Leong JC. Variability of somatosensory-evoked potentials in different stages of scoliosis surgery. *Spine* 1999;24(17):1799-804.

Luk KD, Hu Y, Wong YW, Cheung KM. Evaluation of various evoked potential techniques for spinal cord monitoring during scoliosis surgery. *Spine*. 2001;26(16):1772-7.

Lyon R, Feiner J, Lieberman JA. Progressive suppression of motor evoked potentials during general anesthesia: the phenomenon of "anesthetic fade". *J Neurosurg Anesthesiol*. 2005;17(1):13-9.

MacDonald DB. Individually optimizing posterior tibial somatosensory evoked potential P37 scalp derivations for intraoperative monitoring. *J Clin Neurophysiol* 2001;18(4):364-71.

MacDonald DB, Al-Zayed Z, Dvorak M, et al. Spinal cord monitoring during scoliosis surgery utilizing multiple pulse transcranial electric stimulation motor evoked potentials and somatosensory evoked potentials. *Clinical Neurophysiology* 2001;112 Suppl 1:S100-101.

MacDonald DB, Janusz M. An approach to intraoperative monitoring of thoracoabdominal aneurysm surgery. *J Clin Neurophysiol* 2002;19(1):43-54.

MacDonald DB. Safety of intraoperative transcranial electrical stimulation motor evoked potential monitoring. *J Clin Neurophysiol* 2002;19(5):416-29.

MacDonald DB, Al Zayed Z, Khoudeir I, Stigsby B. Monitoring scoliosis surgery with combined multiple pulse transcranial electric motor and cortical somatosensory-evoked potentials from the lower and upper extremities. *Spine* 2003;28(2):194-203.

MacDonald DB, Stigsby B. Use of D waves in intraoperative monitoring. In: *Proceedings of the Symposium on Intraoperative Neurophysiology* [Ljubjana Institute of Clinical Neurophysiology web site]. Oct 17, 2003;18-28. Available at: www.kclj.si/ikn/Dejavnosti/FAGA/2003/INVI2003.HTM Accessed Nov 12, 2003.

MacDonald DB, Stigsby B, Al-Zayed Z. A comparison between derivation optimization and Cz'-FPz for posterior tibial P37 somatosensory evoked potential intraoperative monitoring. *Clin Neurophysiol* 2004;115:1925-1930.

MacDonald DB, Streletz L, Al-Zayed Z, Abdool S, Stigsby B. Intraoperative neurophysiologic discovery of uncrossed sensory and motor pathways in a patient with horizontal gaze palsy and scoliosis. *Clin Neurophysiol* 2004;115(3):576-582.

MacDonald DB, Al Zayed Z, Stigsby B. Tibial somatosensory evoked potential intraoperative monitoring: Recommendations based on signal to noise ratio analysis of popliteal fossa, optimized P37, standard P37 and P31 potentials. *Clin Neurophysiol* 2005;116(8):1858-69.

MacDonald DB. Safety of transcranial, cortical and subcortical electrical stimulation using different neurophysiological techniques. *Rivista Medica* 2006;12:39-43.

MacDonald DB. Intraoperative motor evoked potential monitoring: Overview and update. *J Clin Monit Comput.* 2006;20(5):347-77. Epub 2006 Jul 11.

MacEwen GD, Bunnell WP, Sriram K. Acute neurological complications in the treatment of scoliosis. A report of the Scoliosis Research Society. *J Bone Joint Surg Am* 1975;57(3):404-8.

Machida M, Weinstein SL, Yamada T, Kimura J, Toriyama S. Dissociation of muscle action potentials and spinal somatosensory evoked potentials after ischemic damage of spinal cord. *Spine.* 1988;13(10):1119-24.

Machida M, Yamada T, Ross M, Kimura J, Hitchon P. Effect of spinal cord ischemia on compound muscle action potentials and spinal evoked potentials following spinal cord stimulation in the dog. *J Spinal Disord.* 1990;3(4):345-52.

Macri S, De Monte A, Greggi T, Parisini P, Zanoni A, Merlini L. Intra-operative spinal cord monitoring in orthopaedics. *Spinal Cord* 2000;38(3):133-9.

Madigan RR, Linton AE, Wallace SL, Dougherty JH, Eisenstadt ML, Powers LB, Zimmerman AW. A new technique to improve cortical-evoked potentials in spinal cord monitoring. A ratio method of analysis. *Spine.* 1987;12(4):330-5.

Marco AP, Rice K. Pseudorhythmia from evoked potential monitoring. *J Neurosurg Anesthesiol.* 2001;13(2):143-5.

Mason DG, Higgins D, Boyd SG, Lloyd-Thomas AR. Effects of isoflurane anaesthesia on the median nerve somatosensory evoked potential in children. *Br J Anaesth* 1992;69(6):562-6.

Matsui Y, Goh K, Shiiya N, Murashita T, Miyama M, Ohba J, Gohda T, Sakuma M, Yasuda K, Tanabe T. Clinical application of evoked spinal cord potentials elicited by direct stimulation of the cord during temporary occlusion of the thoracic aorta. *J Thorac Cardiovasc Surg* 1994;107(6):1519-27.

Matsui Y, Shiiya N, Ishii K, Murashita T, Sasaki S, Sakuma M, Yasuda K. The reliability of evoked spinal cord potentials elicited by direct stimulation of the cord as a monitor of spinal cord ischemia during temporary occlusion of the thoracic aorta. *Panminerva Med* 1997;39(2):78-84.

Maurette P, Simeon F, Castagnera L, Esposito J, Macouillard G, Heraut LA. Propofol anaesthesia alters somatosensory evoked cortical potentials. *Anaesthesia* 1988;43 Suppl:44-5.

May DM, Jones SJ, Crockard HA. Somatosensory evoked potential monitoring in cervical surgery: identification of pre- and intraoperative risk factors associated with neurological deterioration. *J Neurosurg* 1996;85(4):566-73.

McCreery DB, Agnew WF, Yuen TG, Bullara L. Charge density and charge per phase as cofactors in neural injury induced by electrical stimulation. *IEEE Trans Biomed Eng.* 1990;37(10):996-1001.

McCreery DB, Agnew WF, Yuen TG, Bullara LA. Relationship between stimulus amplitude, stimulus frequency and neural damage during electrical stimulation of sciatic nerve of cat. *Med Biol Eng Comput.* 1995;33(3 Spec No):426-9.

McPherson RW, Mahla M, Johnson R, Traystman RJ. Effects of enflurane, isoflurane, and nitrous oxide on somatosensory evoked potentials during fentanyl anesthesia. *Anesthesiology* 1985;62(5):626-33.

McPherson RW, Sell B, Traystman RJ. Effects of thiopental, fentanyl, and etomidate on upper extremity somatosensory evoked potentials in humans. *Anesthesiology* 1986;65(6):584-9.

Merton PA and Morton HB. Stimulation of the cerebral cortex in the intact human subject. *Nature* 1980;285:227.

Meylaerts SA, Jacobs MJ, van Iterson V, De Haan P, Kalkman CJ. Comparison of transcranial motor evoked potentials and somatosensory evoked potentials during thoracoabdominal aortic aneurysm repair. *Ann Surg* 1999;230(6):742-9.

Meylaerts SA, De Haan P, Kalkman CJ, Lips J, De Mol BA, Jacobs MJ. The influence of regional spinal cord hypothermia on transcranial myogenic motor-evoked potential monitoring and the efficacy of spinal cord ischemia detection. *J Thorac Cardiovasc Surg* 1999;118(6):1038-45.

Meylaerts SA, De Haan P, Kalkman CJ, Jaspers J, Vanicky I, Jacobs MJ. Prevention of paraplegia in pigs by selective segmental artery perfusion during aortic cross-clamping. *J Vasc Surg* 2000;32(1):160-70.

Michel F, Rubini J, Grand C, Berard J, Kohler R, Michel CR. Neurological complications of surgery for spinal deformities. *Rev Chir Orthop Reparatrice Appar Mot* 1992;78(2):90-100.

Michenfelder JD. Intraoperative monitoring of sensory evoked potentials may be neither a proven nor an indicated technique. *J Clin Monit* 1987;3(1):45-7.

Minahan RE, Sepkuty JP, Lesser RP, Sponseller PD, Kostuik JP. Anterior spinal cord injury with preserved neurogenic 'motor' evoked potentials. *Clin Neurophysiol* 2001;112(8):1442-50.

Mineiro J, Weinstein SL. Delayed postoperative paraparesis in scoliosis surgery. A case report. *Spine* 1997;22(14):1668-72.

Mineiro J, Weinstein SL. Delayed postoperative paraparesis in scoliosis surgery. A case report. *Spine*. 1997;22(14):1668-72.

Miura T, Sonoo M, Shimizu T. Establishment of standard values for the latency, interval and amplitude parameters of tibial nerve somatosensory evoked potentials (SEPs). *Clin Neurophysiol*. 2003;114(7):1367-78.

Mizrahi EM, Crawford ES. Somatosensory evoked potentials during reversible spinal cord ischemia in man. *Electroencephalogr Clin Neurophysiol* 1984;58(2):120-6.

Mochida K, Komori H, Okawa A, Shinomiya K. Evaluation of motor function during thoracic and thoracolumbar spinal surgery based on motor-evoked potentials using train spinal stimulation. *Spine* 1997;22(12):1385-93.

Moe JH. Complications of scoliosis treatment. *Clin Orthop* 1967;53:21-30.

More RC, Nuwer MR, Dawson EG. Cortical evoked potential monitoring during spinal surgery: sensitivity, specificity, reliability, and criteria for alarm. *J Spinal Disord* 1988;1(1):75-80.

Morota N, Deletis V, Constantini S, Kofler M, Cohen H, Epstein FJ. The role of motor evoked potentials during surgery for intramedullary spinal cord tumors. *Neurosurgery* 1997;41(6):1327-36.

Mostegl A, Bauer R. The application of somatosensory-evoked potentials in orthopedic spine surgery. *Arch Orthop Trauma Surg* 1984;103(3):179-84.

Mostegl A, Bauer R, Eichenauer M. Intraoperative somatosensory potential monitoring. A clinical analysis of 127 surgical procedures. *Spine* 1988;13(4):396-400.

Murakami H, Tsukube T, Kawanishi Y, Okita Y. Transcranial myogenic motor-evoked potentials after transient spinal cord ischemia predicts neurologic outcome in rabbits. *J Vasc Surg*. 2004;39(1):207-13.

Mustain WD, Kendig RJ. Dissociation of neurogenic motor and somatosensory evoked potentials. A case report. *Spine* 1991;16(7):851-3.

Nagle KJ, Emerson RG, Adams DC, Heyer EJ, Roye DP, Schwab FJ, Weidenbaum M, McCormick P, Pile-Spellman J, Stein BM, Farcy JP, Gallo EJ, Dowling KC, Turner CA. Intraoperative monitoring of motor evoked potentials: a review of 116 cases. *Neurology* 1996;47(4):999-1004.

Naito M, Owen JH, Bridwell KH, Sugioka Y. Effects of distraction on physiologic integrity of the spinal cord, spinal cord blood flow, and clinical status. *Spine* 1992;17(10):1154-8.

Nakao Y, Piccirillo E, Falcioni M, Taibah A, Kobayashi T, Sanna M. Electromyographic evaluation of facial nerve damage in acoustic neuroma surgery. *Otol Neurotol* 2001;22(4):554-7.

Niimi Y, Sala F, Deletis V, Setton A, de Camargo AB, Berenstein A. Neurophysiologic monitoring and pharmacologic provocative testing for embolization of spinal cord arteriovenous malformations. *Am J Neuroradiol.* 2004;25(7):1131-8.

Niznik G, Transfeldt EE, Shichijo F, Ohshima T, Pomeranz B. Spinal sensory and motor tract activation after epidural electrical stimulation in the cat. *Spine* 1990;15(7):623-9.

Noel P, Deltenre P, Lamoureux J, Capouet V, de Ville A, Govaerts MJ. Neurophysiologic detection of a unilateral motor deficit occurring during the noncritical phase of scoliosis surgery. *Spine* 1994;19(21):2399-402.

Noonan KJ, Walker T, Feinberg JR, Nagel M, Didelot W, Lindseth R. Factors related to false- versus true-positive neuromonitoring changes in adolescent idiopathic scoliosis surgery. *Spine.* 2002;27(8):825-30.

Norcross-Nechay K, Mathew T, Simmons JW, Hadjipavlou A. Intraoperative somatosensory evoked potential findings in acute and chronic spinal canal compromise. *Spine* 1999;24(10):1029-33.

North RB, Drenger B, Beattie C, McPherson RW, Parker S, Reitz BA, Williams GM. Monitoring of spinal cord stimulation evoked potentials during thoracoabdominal aneurysm surgery. *Neurosurgery* 1991;28(2):325-30.

Novak K, de Camargo AB, Neuwirth M, Kothbauer K, Amassian VE, Deletis V. The refractory period of fast conducting corticospinal tract axons in man and its implications for intraoperative monitoring of motor evoked potentials. *Clin Neurophysiol.* 2004;115(8):1931-41.

Nuwer MR, Dawson EC. Intraoperative evoked potential monitoring of the spinal cord. A restricted filter, scalp method during Harrington instrumentation for scoliosis. *Clin Orthop* 1984;(183):42-50.

Nuwer MR. Electrophysiologic evaluation and monitoring of spinal cord and root function. *Neurosurg Clin N Am* 1990;1(3):533-49.

Nuwer MR, Dawson EG, Carlson LG, Kanim LE, Sherman JE. Somatosensory evoked potential spinal cord monitoring reduces neurologic deficits after scoliosis surgery: results of a large multicenter survey. *Electroencephalogr Clin Neurophysiol* 1995;96(1):6-11.

Nuwer MR. Spinal cord monitoring with somatosensory techniques. *J Clin Neurophysiol* 1998;15:183-93.

Nuwer MR. Spinal cord monitoring. *Muscle Nerve* 1999;22(12):1620-30.

O'Brien MF, Lenke LG, Bridwell KH, Padberg A, Stokes M. Evoked potential monitoring of the upper extremities during thoracic and lumbar spinal deformity surgery: a prospective study. *J Spinal Disord* 1994;7(4):277-84.

Oberle J, Antoniadis G, Rath SA, Richter HP. Intra-operative electrophysiological diagnosis of spinal root avulsion during surgical repair of brachial plexus stretch injuries. *Acta Neurochir (Wien)* 1997;139(3):238-9.

Okada K, Sueda T, Shikata H, Orihashi K, Morita S, Hirai S, Suehiro M, Wada H, Matsuura Y. Evoked spinal cord potential monitoring reveals peroneal nerve ischemia during thoracoabdominal repair: a case report. *Ann Thorac Cardiovasc Surg*. 1999;5(5):350-2.

Okamoto Y, Murakami M, Nakagawa T, Murata A, Moriya H. Intraoperative spinal cord monitoring during surgery for aortic aneurysm: application of spinal cord evoked potential. *Electroencephalogr Clin Neurophysiol* 1992;84(4):315-20.

Owen JH, Laschinger J, Bridwell K, Shimon S, Nielsen C, Dunlap J, Kain C. Sensitivity and specificity of somatosensory and neurogenic-motor evoked potentials in animals and humans. *Spine* 1988;13(10):1111-8.

Owen JH, Jenny AB, Naito M, Weber K, Bridwell KH, McGhee R. Effects of spinal cord lesioning on somatosensory and neurogenic-motor evoked potentials. *Spine* 1989;14(7):673-82.

Owen JH, Naito M, Bridwell KH. Relationship among level of distraction, evoked potentials, spinal cord ischemia and integrity, and clinical status in animals. *Spine* 1990;15(9):852-7.

Owen JH, Naito M, Bridwell KH, Oakley DM. Relationship between duration of spinal cord ischemia and postoperative neurologic deficits in animals. *Spine* 1990;15(9):846-51.

Owen JH, Naito M, Bridwell KH, Oakley DM. Relationship between duration of spinal cord ischemia and postoperative neurologic deficits in animals. *Spine* 1990;15(7):618-22.

Owen JH, Bridwell KH, Grubb R, Jenny A, Allen B, Padberg AM, Shimon SM. The clinical application of neurogenic motor evoked potentials to monitor spinal cord function during surgery. *Spine* 1991;16(8 Suppl):S385-90.

Owen JH, Bridwell KH, Lenke LG. Innervation pattern of dorsal roots and their effects on the specificity of dermatomal somatosensory evoked potentials. *Spine* 1993;18(6):748-54.

Owen JH. Intraoperative stimulation of the spinal cord for prevention of spinal cord injury. *Adv Neurol* 1993;63:271-88.

Owen JH, Lenke LG, Danto J. Spinal cord monitoring in patients undergoing surgery for neuromuscular scoliosis. *Spine* 1994;19(20):2371-2.

Owen JH, Sponseller PD, Szymanski J, Hurdle M. Efficacy of multimodality spinal cord monitoring during surgery for neuromuscular scoliosis. *Spine* 1995;20(13):1480-8.

Owen JH. The application of intraoperative monitoring during surgery for spinal deformity. *Spine* 1999;24(24):2649-62.

Padberg AM, Russo MH, Lenke LG, Bridwell KH, Komanetsky RM. Validity and reliability of spinal cord monitoring in neuromuscular spinal deformity surgery. *J Spinal Disord* 1996;9(2):150-8.

Padberg AM, Wilson-Holden TJ, Lenke LG, Bridwell KH. Somatosensory- and motor-evoked potential monitoring without a wake-up test during idiopathic scoliosis surgery. An accepted standard of care. *Spine* 1998;23(12):1392-400.

Padberg AM, Bridwell KH. Spinal cord monitoring: current state of the art. *Orthop Clin North Am* 1999;30(3):407-33.

Paisansathan C, Koenig HM, Wheeler PJ, Baughman VL, Hoffman WE. Loss of SSEP during sitting craniotomy. *J Neurosurg Anesthesiol*. 2003;15(4):327-9.

Papastefanou SL, Henderson LM, Smith NJ, Hamilton A, Webb JK. Surface electrode somatosensory-evoked potentials in spinal surgery: implications for indications and practice. *Spine* 2000;25(19):2467-72.

Partanen J, Merikanto J, Kokki H, Kilpelainen R, Koistinen A. Antidromic corticospinal tract potential of the brain. *Clin Neurophysiol*. 2000;111(3):489-95.

Passuti N, Delecrin J, Pereon Y, Nguyen ST. Neurologic complications of surgical treatments. *Bull Acad Natl Med* 1999;183(4):769-73.

Pathak KS, Amaddio MD, Scoles PV, Shaffer JW, Mackay W. Effects of halothane, enflurane, and isoflurane in nitrous oxide on multilevel somatosensory evoked potentials. *Anesthesiology* 1989;70(2):207-12.

Patton HD, Amassian VE. Single- and multiple-unit analysis of cortical stage of pyramidal tract activation. *J Neurophysiol* 1954;17:345-63.

Pechstein U, Cedzich C, Nadstawek J, Schramm J. Transcranial high-frequency repetitive electrical stimulation for recording myogenic motor evoked potentials with the patient under general anesthesia. *Neurosurgery* 1996;39(2):335-43; discussion 343-4.

Pechstein U, Nadstawek J, Zentner J, Schramm J. Isoflurane plus nitrous oxide versus propofol for recording of motor evoked potentials after high frequency repetitive electrical stimulation. *Electroencephalogr Clin Neurophysiol* 1998;108(2):175-81.

Pelosi L, Jardine A, Webb JK. Neurological complications of anterior spinal surgery for kyphosis with normal somatosensory evoked potentials (SEPs). *J Neurol Neurosurg Psychiatry* 1999;66(5):662-4.

Pelosi L, Stevenson M, Hobbs GJ, Jardine A, Webb JK. Intraoperative motor evoked potentials to transcranial electrical stimulation during two anaesthetic regimens. *Clin Neurophysiol* 2001;112(6):1076-87.

Pelosi L, Lamb J, Grevitt M, Mehdian SMH, Webb JK, Blumhardt LD. Combined monitoring of motor and somatosensory evoked potentials in orthopaedic spinal surgery. *Clinical Neurophysiology* 2002;113:1082–1091.

Pereon Y, Bernard JM. Intraoperative monitoring of motor evoked potentials. *Anesth Analg* 1994;79(6):1206.

Pereon Y, Bernard JM, Fayet G, Delecrin J, Passuti N, Guiheneuc P. Usefulness of neurogenic motor evoked potentials for spinal cord monitoring: findings in 112 consecutive patients undergoing surgery for spinal deformity. *Electroencephalogr Clin Neurophysiol* 1998;108(1):17-23.

Pereon Y, Nguyen The Tich S. Monitoring the neurogenic mixed evoked potentials during surgery of the spine: a good solution? *Neurophysiol Clin* 1998;28(4):321-34.

Pereon Y, Nguyen The Tich S, Delecrin J, Passuti N. Somatosensory- and motor-evoked potential monitoring without a wake-up test during idiopathic scoliosis surgery: an accepted standard of care. *Spine* 1999;24(11):1169-70.

Pereon Y, Delecrin J, Nguyen The Tich SN, Bertrand-Vasseur A, Passuti N. Successful monitoring of neurogenic mixed evoked potentials elicited by anterior spinal cord stimulation through thoracoscopy during spine surgery. *Spine* 1999;24(19):2025-9.

Pereon Y, Bernard JM, Nguyen The Tich S, Genet R, Petitfaux F, Guiheneuc P. The effects of desflurane on the nervous system: from spinal cord to muscles. *Anesth Analg* 1999;89(2):490-5.

Perlik SJ, VanEgeren R, Fisher MA. Somatosensory evoked potential surgical monitoring. Observations during combined isoflurane-nitrous oxide anesthesia. *Spine* 1992;17(3):273-6.

Peterson DO, Drummond JC, Todd MM. Effects of halothane, enflurane, isoflurane, and nitrous oxide on somatosensory evoked potentials in humans. *Anesthesiology* 1986;65(1):35-40.

Phillips LH 2nd, Blanco JS, Sussman MD. Direct spinal stimulation for intraoperative monitoring during scoliosis surgery. *Muscle Nerve* 1995;18(3):319-25.

Piatt JH Jr, Radtke RA, Erwin CW. Limitations of brain stem auditory evoked potentials for intraoperative monitoring during a posterior fossa operation: case report and technical note. *Neurosurgery* 1985;16(6):818-21.

Pollock JC, Jamieson MP, McWilliam R. Somatosensory evoked potentials in the detection of spinal cord ischemia in aortic coarctation repair. *Ann Thorac Surg*. 1986;41(3):251-4.

Porkkala T, Jantti V, Kaukinen S, Hakkinen V. Somatosensory evoked potentials during isoflurane anaesthesia. *Acta Anaesthesiol Scand* 1994;38(3):206-10.

Porkkala T, Jantti V, Kaukinen S, Hakkinen V. Nitrous oxide has different effects on the EEG and somatosensory evoked potentials during isoflurane anaesthesia in patients. *Acta Anaesthesiol Scand* 1997;41(4):497-501.

Porkkala T, Kaukinen S, Hakkinen V, Jantti V. Median nerve somatosensory evoked potentials during isoflurane anaesthesia. *Can J Anaesth* 1997;44(9):963-8.

Prestor B, Zgur T, Dolenc VV. Epidural and subpial cortico-spinal potentials evoked by transcutaneous motor cortex stimulation during spinal cord surgery. *Electroencephalogr Clin Neurophysiol Suppl*. 1990;41:348-51.

Prokop A, Meyer GP, Walter M, Erasmi H. Validity of SEP monitoring in carotid surgery. Review and own results. *J Cardiovasc Surg (Torino)*. 1996;37(4):337-42.

Qayumi KA, Janusz MT, Jamieson EW, Chow VD, Dry GM. Transcranial magnetic stimulation: use of motor evoked potentials in the evaluation of surgically induced spinal cord ischemia. *J Spinal Cord Med* 1997;20(4):395-401.

Quinones-Hinojosa A, Gulati M, Lyon R, Gupta N, Yingling C. Spinal cord mapping as an adjunct for resection of intramedullary tumors: surgical technique with case illustrations. *Neurosurgery*. 2002;51(5):1199-206; discussion 1206-7.

Quinones-Hinojosa A, Gadkary CA, Gulati M, von Koch CS, Lyon R, Weinstein PR, Yingling CD. Neurophysiological monitoring for safe surgical tethered cord syndrome release in adults. *Surg Neurol*. 2004;62(2):127-33.

Quiñones-Hinojosa A, Alam M, Lyon R, Yingling CD, Lawton MT. Transcranial motor evoked potentials during basilar artery aneurysm surgery: technique application for 30 consecutive patients. *Neurosurgery*. 2004;54(4):916-24; discussion 924.

Quiñones-Hinojosa A, Lyon R, Zada G, Lamborn KR, Gupta N, Parsa AT, McDermott MW, Weinstein PR. Changes in transcranial motor evoked potentials during

intramedullary spinal cord tumor resection correlate with postoperative motor function. *Neurosurgery*. 2005;56(5):982-93.

Raudzens PA. Intraoperative monitoring of evoked potentials. *Ann N Y Acad Sci* 1982;388:308-26.

Raynor BL, Lenke LG, Kim Y, Hanson DS, Wilson-Holden TJ, Bridwell KH, Padberg AM. Can triggered electromyograph thresholds predict safe thoracic pedicle screw placement? *Spine*. 2002;27(18):2030-5.

Rehberg B, Grünewald M, Baars J, Fuegener K, Urban BW, Kox WJ. Monitoring of immobility to noxious stimulation during sevoflurane anesthesia using the spinal H-reflex. *Anesthesiology*. 2004;100(1):44-50.

Reuter DG, Tacker WA Jr, Badylak SF, Voorhees WD 3rd, Konrad PE. Correlation of motor-evoked potential response to ischemic spinal cord damage. *J Thorac Cardiovasc Surg* 1992;104(2):262-72.

Rittmeister M, Leyendecker K, Kurth A, Schmitt E. Cauda equina compression due to a laminar hook: A late complication of posterior instrumentation in scoliosis surgery. *Eur Spine J* 1999;8(5):417-20.

Robertazzi RR, Cunningham JN Jr. Intraoperative adjuncts of spinal cord protection. *Semin Thorac Cardiovasc Surg* 1998;10(1):29-34.

Robertazzi RR, Cunningham JN Jr. Monitoring of somatosensory evoked potentials: a primer on the intraoperative detection of spinal cord ischemia during aortic reconstructive surgery. *Semin Thorac Cardiovasc Surg* 1998;10(1):11-7.

Rodi Z, Deletis V, Morota N, Vodusek DB. Motor evoked potentials during brain surgery. *Pflugers Arch* 1996;431(6 Suppl 2):R291-2.

Rodi Z, Vodusek DB. Intraoperative monitoring of the bulbocavernosus reflex: the method and its problems. *Clin Neurophysiol* 2001;112(5):879-83.

Rodi Z, Straus I, Denic´ K, Deletis V, Vodusek DB. Transient paraplegia revealed by intraoperative neurophysiological monitoring: was it caused by the epidural anesthetic or an epidural hematoma? *Anesth Analg*. 2003;96(6):1785-8.

Romstock J, Strauss C, Fahlbusch R. Continuous electromyography monitoring of motor cranial nerves during cerebellopontine angle surgery. *J Neurosurg* 2000;93(4):586-93.

Rose RD. Sensory component of cervically evoked motor potentials. *Med Hypotheses* 1996;46(6):577-9.

Rose RD, Welch WC, Balzer JR, Jacobs GB. Persistently electrified pedicle stimulation instruments in spinal instrumentation. Technique and protocol development. *Spine* 1997;22(3):334-43.

Rose RD. Removing the antidromically driven sensory component from cervically evoked motor potentials. *Med Hypotheses* 1998;50(2):147-54.

Rothwell J, Burke D, Hicks R, Stephen J, Woodforth I, Crawford M. Transcranial electrical stimulation of the motor cortex in man: further evidence for the site of activation. *J Physiol.* 1994;481 (Pt 1):243-50.

Roy EP 3rd, Gutmann L, Riggs JE, Jones ET, Byrd JA, Ringel RA. Intraoperative somatosensory evoked potential monitoring in scoliosis. *Clin Orthop* 1988;(229):94-8.

Ruskin KJ, Deletis V, Morota N. Anesthesia and intraoperative monitoring. *J Neurosurg* 1994;80(5):946-7.

Russ W, Kling D, Schwandt HJ, Krumholz W, Hempelmann G. Effects of thiopentone and etomidate on median nerve somatosensory evoked responses. *Neurol Res* 1989;11(1):51-6.

Ryan TP, Britt RH. Spinal and cortical somatosensory evoked potential monitoring during corrective spinal surgery with 108 patients. *Spine.* 1986;11(4):352-61.

Rytky S, Huotari AM, Alahuhta S, Remes R, Suominen K, Jantti V. Tibial nerve somatosensory evoked potentials during EEG suppression in sevoflurane anaesthesia. *Clin Neurophysiol* 1999;110(9):1655-8.

Sakuma J, Suzuki K, Sasaki T, Matsumoto M, Oinuma M, Kawakami M, Itakura T, Kodama N. Monitoring and preventing blood flow insufficiency due to clip rotation after the treatment of internal carotid artery aneurysms. *J Neurosurg.* 2004;100(5):960-2.

Sala F, Niimi Y, Berenstein A, Deletis V. Neuroprotective role of neurophysiological monitoring during endovascular procedures in the spinal cord. *Ann N Y Acad Sci* 2001;939:126-36.

Sala F, Lanteri P. Brain surgery in motor areas: the invaluable assistance of intraoperative neurophysiological monitoring. *J Neurosurg Sci.* 2003;47(2):79-88.

Sala F, Lanteri P, Bricolo A. Motor evoked potential monitoring for spinal cord and brain stem surgery. *Adv Tech Stand Neurosurg.* 2004;29:133-69.

Salvian AJ, Taylor DC, Hsiang YN, Hildebrand HD, Litherland HK, Humer MF, Teal PA, MacDonald DB. Selective shunting with EEG monitoring is safer than routine shunting for carotid endarterectomy. *Cardiovasc Surg* 1997;5(5):481-5.

Samra SK, Dy EA, Welch KB, Lovely LK, Graziano GP. Remifentanyl- and fentanyl-based anesthesia for intraoperative monitoring of somatosensory evoked potentials. *Anesth Analg.* 2001;92(6):1510-5.

Sartorius CJ, Wright G. Intraoperative brain mapping in a community setting--technical considerations. *Surg Neurol.* 1997;47(4):380-8.

Schepens M, Dossche K, Morshuis W, Heijmen R, van Dongen E, Ter Beek H, Kelder H, Boezeman E. Introduction of adjuncts and their influence on changing results in 402 consecutive thoracoabdominal aortic aneurysm repairs. *Eur J Cardiothorac Surg*. 2004;25(5):701-7.

Schepens MA, Boezeman EH, Hamerlijnck RP, ter Beek H, Vermeulen FE. Somatosensory evoked potentials during exclusion and reperfusion of critical aortic segments in thoracoabdominal aortic aneurysm surgery. *J Card Surg* 1994;9(6):692-702.

Scheufler KM, Zentner J. Total intravenous anesthesia for intraoperative monitoring of the motor pathways: an integral view combining clinical and experimental data. *J Neurosurg*. 2002;96(3):571-9.

Scheufler KM, Thees C, Nadstawek J, Zentner J. S(+)-ketamine attenuates myogenic motor-evoked potentials at or distal to the spinal alpha-motoneuron. *Anesth Analg*. 2003;96(1):238-44.

Scheufler KM, Reinacher PC, Blumrich W, Zentner J, Priebe HJ. The Modifying Effects of Stimulation Pattern and Propofol Plasma Concentration on Motor-Evoked Potentials. *Anesth Analg* 2005;100:440-7.

Schindler E, Muller M, Zickmann B, Osmer C, Wozniak G, Hempelmann G. Modulation of somatosensory evoked potentials under various concentrations of desflurane with and without nitrous oxide. *J Neurosurg Anesthesiol* 1998;10(4):218-23.

Schlake HP, Goldbrunner RH, Milewski C, Krauss J, Trautner H, Behr R, Sorensen N, Helms J, Roosen K. Intra-operative electromyographic monitoring of the lower cranial motor nerves (LCN IX-XII) in skull base surgery. *Clin Neurol Neurosurg*. 2001;103(2):72-82.

Schmidt J, Hering W, Albrecht S. Total intravenous anesthesia with propofol and remifentanyl. Results of a multicenter study of 6,161 patients. *Anaesthesist*. 2005;54(1):17-28.

Schramm J, Kurthen M. Recent developments in neurosurgical spinal cord monitoring. *Paraplegia* 1992;30(9):609-16.

Schubert A, Drummond JC, Peterson DO, Saidman LJ. The effect of high-dose fentanyl on human median nerve somatosensory-evoked responses. *Can J Anaesth* 1987;34(1):35-40.

Schubert A, Todd MM, Luerssen TG, Hicks GE. Loss of intraoperative evoked responses during dorsal column surgery associated with isolated postoperative sensory deficit. *J Clin Monit* 1987;3(4):277-81.

Schubert A, Licina MG, Lineberry PJ. The effect of ketamine on human somatosensory evoked potentials and its modification by nitrous oxide [published erratum appears in *Anesthesiology* 1990;72(6):1104] *Anesthesiology* 1990;72(1):33-9.

Schwartz DM, Drummond DS, Ecker ML. Influence of rigid spinal instrumentation on the neurogenic motor evoked potential. *J Spinal Disord* 1996;9(5):439-45.

Schwartz DM, Schwartz JA, Pratt RE Jr, Wierzbowski LR, Sestokas AK. Influence of nitrous oxide on posterior tibial nerve cortical somatosensory evoked potentials. *J Spinal Disord* 1997;10(1):80-6.

Schwartz DM, Drummond DS, Hahn M, Ecker ML, Dormans JP. Prevention of positional brachial plexopathy during surgical correction of scoliosis. *J Spinal Disord* 2000;13(2):178-82.

Sebel PS, Erwin CW, Neville WK. Effects of halothane and enflurane on far and near field somatosensory evoked potentials. *Br J Anaesth* 1987;59(12):1492-6.

Selesnick SH. Optimal stimulus duration for intraoperative facial nerve monitoring. *Laryngoscope* 1999;109(9):1376-85.

Shahin GM, Hamerlijnck RP, Schepens MA, Ter Beek HT, Vermeulen FE, Boezeman EH. Upper and lower extremity somatosensory evoked potential recording during surgery for aneurysms of the descending thoracic aorta. *Eur J Cardiothorac Surg* 1996;10(5):299-304.

Shimoda T, Inoue S, Kakimoto M, Sakamoto T, Kawaguchi M, Kitaguchi K, Furuya H, Tabayashi N, Taniguchi S. Intraoperative unilateral changes in myogenic motor evoked potentials in patients undergoing thoracic aortic surgery. *Masui*. 2001;50(8):874-7.

Shokoku S, Uchida H, Teramoto S. An experimental study on spinal cord ischemia during cross-clamping of the thoracic aorta: the monitoring of spinal cord ischemia with motor evoked potential by transcranial stimulation of the cerebral cortex in dogs. *Surg Today* 1993;23(12):1068-73.

Sihle-Wissel M, Scholz M, Cunitz G. Transcranial magnetic-evoked potentials under total intravenous anaesthesia and nitrous oxide. *Br J Anaesth* 2000;85(3):465-7.

Slimp JC, Stolov WC, Wagner TA. Spine and scalp recordings as a function of intensity. A model for changes during spinal cord monitoring. *Spine*. 1996;21(1):99-103.

Sloan TB, Koht A. Depression of cortical somatosensory evoked potentials by nitrous oxide. *Br J Anaesth* 1985;57(9):849-52.

Sloan TB, Ronai AK, Toleikis JR, Koht A. Improvement of intraoperative somatosensory evoked potentials by etomidate. *Anesth Analg* 1988;67(6):582-5.

Sloan TB, Heyer EJ. Anesthesia for Intraoperative Neurophysiologic Monitoring of the Spinal Cord. *J Clin Neurophysiol* 2002;19:430-443.

Sloan TB. Electrophysiologic monitoring during surgery to repair the thoracoabdominal aorta. *Semin Cardiothorac Vasc Anesth*. 2004;8(2):113-25.

Sobottka SB, Schackert G, May SA, Wiegler M, Reiss G. Intraoperative facial nerve monitoring (IFNM) predicts facial nerve outcome after resection of vestibular schwannoma. *Acta Neurochir (Wien)* 1998;140(3):235-42; discussion 242-3.

Spielholz NI, Benjamin MV, Engler GL, Ransohoff J. Somatosensory evoked potentials during decompression and stabilization of the spine. *Methods and findings. Spine* 1979;4(6):500-5.

Stecker MM, Cheung AT, Pochettino A, Kent GP, Patterson T, Weiss SJ, Bavaria JE. Deep hypothermic circulatory arrest: I. Effects of cooling on electroencephalogram and evoked potentials. *Ann Thorac Surg.* 2001;71(1):14-21.

Stecker MM, Escherich A, Patterson T, Bavaria JE, Cheung AT. Effects of acute hypoxemia/ischemia on EEG and evoked responses at normothermia and hypothermia in humans. *Med Sci Monit.* 2002;8(4):CR223-8.

Stecker MM. Transcranial electric stimulation of motor pathways: a theoretical analysis. *Comput Biol Med* 2005;35(2):133-55.

Stephen JP, Sullivan MR, Hicks RG, Burke DJ, Woodforth IJ, Crawford MR. Cotrel-dubousset instrumentation in children using simultaneous motor and somatosensory evoked potential monitoring. *Spine* 1996;21(21):2450-7.

Stinson LW Jr, Murray MJ, Jones KA, Assef SJ, Burke MJ, Behrens TL, Lennon RL. A computer-controlled, closed-loop infusion system for infusing muscle relaxants: its use during motor-evoked potential monitoring. *J Cardiothorac Vasc Anesth* 1994;8(1):40-4.

Strahm C, Min K, Boos N, Ruetsch Y, Curt A. Reliability of perioperative SSEP recordings in spine surgery. *Spinal Cord.* 2003;41(9):483-9.

Stühmeier KD, Grabitz K, Mainzer B, Sandmann W, Tarnow J. Use of the electrospinogram for predicting harmful spinal cord ischemia during repair of thoracic or thoracoabdominal aortic aneurysms. *Anesthesiology.* 1993;79(6):1170-6; discussion 27A-28A.

Su CF, Haghghi SS, Oro JJ, Gaines RW. "Backfiring" in spinal cord monitoring. *Spine* 1992;17:504-508.

Sueda T, Okada K, Morita S, Wada H, Watari M, Orihashi K, Shikata H, Matsuura Y. Three different approaches to spinal cord monitoring for the prediction of spinal cord ischemia during thoracic aortic aneurysm surgery. *Ann Thorac Cardiovasc Surg* 1998;4(5):290-3.

Sueda T, Okada K, Watari M, Orihashi K, Shikata H, Matsuura Y. Evaluation of motor- and sensory-evoked potentials for spinal cord monitoring during thoracoabdominal aortic aneurysm surgery. *Jpn J Thorac Cardiovasc Surg* 2000;48(1):60-5.

Sueda T, Morita S, Okada K, Orihashi K, Shikata H, Matsuura Y. Selective intercostal arterial perfusion during thoracoabdominal aortic aneurysm surgery. *Ann Thorac Surg.* 2000;70(1):44-7.

Suihko V, Eskola HJ. Transcranial electrical stimulator producing high amplitude pulses and pulse trains. *Med Eng Technol* 1998;22(5):211-5.

Suzuki K, Kodama N, Sasaki T, Matsumoto M, Konno Y, Sakuma J, Oinuma M, Murakawa M. Intraoperative monitoring of blood flow insufficiency in the anterior choroidal artery during aneurysm surgery. *J Neurosurg.* 2003;98(3):507-14.

Svensson LG, Patel V, Robinson MF, Ueda T, Roehm JO Jr, Crawford ES. Influence of preservation or perfusion of intraoperatively identified spinal cord blood supply on spinal motor evoked potentials and paraplegia after aortic surgery. *J Vasc Surg.* 1991;13(3):355-65.

Svensson LG, Crawford ES, Hess KR, Coselli JS, Safi HJ. Experience with 1509 patients undergoing thoracoabdominal aortic operations. *J Vasc Surg* 1993;17(2):357-68; discussion 368-70.

Szelenyi A, Bueno De Camargo A, Deletis V. Neurophysiological evaluation of the corticospinal tract by D-wave recordings in young children. *Childs Nerv Syst.* 2003;19(1):30-4.

Szelenyi A, Kothbauer K, de Camargo AB, Langer D, Flamm ES, Deletis V. Motor evoked potential monitoring during cerebral aneurysm surgery: technical aspects and comparison of transcranial and direct cortical stimulation. *Neurosurgery.* 2005;57(4 Suppl):331-8.

Szelényi A, Bueno de Camargo A, Flamm E, Deletis V. Neurophysiological criteria for intraoperative prediction of pure motor hemiplegia during aneurysm surgery. Case report. *J Neurosurg.* 2003;99(3):575-8.

Tabaraud F, Boulesteix JM, Moulies D, Longis B, Lansade A, Terrier G, Vallat JM, Dumas M, Hugont J. Monitoring of the motor pathway during spinal surgery. *Spine* 1993;18(5):546-50.

Tanaka S, Kobayashi I, Sagiuchi T, Takanashi J, Iwamoto K, Sato S, Fujii K. Compensation of intraoperative transcranial motor-evoked potential monitoring by compound muscle action potential after peripheral nerve stimulation. *J Clin Neurophysiol.* 2005;22(4):271-4.

Taniguchi M, Schramm J. Motor evoked potentials facilitated by an additional peripheral nerve stimulation. *Electroencephalogr Clin Neurophysiol Suppl* 1991;43:202-11.

Taniguchi M, Nadstawek J, Pechstein U, Schramm J. Total intravenous anesthesia for improvement of intraoperative monitoring of somatosensory evoked potentials during aneurysm surgery. *Neurosurgery* 1992;31(5):891-7; discussion 897.

Taniguchi M, Cedzich C, Schramm J. Modification of cortical stimulation for motor evoked potentials under general anesthesia: technical description. *Neurosurgery* 1993;32(2):219-26.

Taniguchi M, Nadstawek J, Langenbach U, Bremer F, Schramm J. Effects of four intravenous anesthetic agents on motor evoked potentials elicited by magnetic transcranial stimulation. *Neurosurgery* 1993;33(3):407-15; discussion 415.

Taunt CJ, Sidhu KS, Andrew SA. Somatosensory Evoked Potential Monitoring During Anterior Cervical Discectomy and Fusion. *Spine* 2005;30(17):1970-1972.

Taylor BA, Fennelly ME, Taylor A, Farrell J. Temporal summation--the key to motor evoked potential spinal cord monitoring in humans. *J Neurol Neurosurg Psychiatry* 1993;56(1):104-6.

Taylor BA, Webb PJ, Hetreed M, Mulukutla RD, Farrell J. Delayed postoperative paraplegia with hypotension in adult revision scoliosis surgery. *Spine* 1994;19(4):470-4.

Terrier G, Lansade A, Preux PM, Tabaraud F, Moulies D. Effects of isoflurane and halothane on motor evoked potentials in the rabbit. *Ann Fr Anesth Reanim* 1993;12(3):284-8.

Thees C, Scheufler KM, Nadstawek J, Pechstein U, Hanisch M, Juntke R, Zentner J, Hoeft A. Influence of fentanyl, alfentanil, and sufentanil on motor evoked potentials. *J Neurosurg Anesthesiol* 1999;11(2):112-8.

Thompson PD, Day BL, Crockard HA, Calder I, Murray NM, Rothwell JC, Marsden CD. Intra-operative recording of motor tract potentials at the cervico-medullary junction following scalp electrical and magnetic stimulation of the motor cortex. *J Neurol Neurosurg Psychiatry* 1991;54(7):618-23.

Thuet ED, Padberg AM, Raynor BL, Bridwell KH, Riew KD, Taylor BA, Lenke LG. Increased Risk of Postoperative Neurologic Deficit for Spinal Surgery Patients With Unobtainable Intraoperative Evoked Potential Data. *Spine* 2005; 30(18):2094-2103.

Tinazzi M, Zanette G, Fiaschi A, Maguiere F. Effects of stimulus rate on the cortical posterior tibial nerve SEPs: A topographic study. *Electroencephalogr Clin Neurophysiol* 1996;100:210-19.

Tohdoh Y, Sumita S, Kawamata T, Omote K, Kawana S, Namiki A. Acute respiratory and metabolic acidosis induced by excessive muscle contraction during spinal evoked stimulation. *Br J Anaesth*. 2001;86(4):589-93.

Tokimura H, Yamagami M, Tokimura Y, Asakura T, Atsuchi M. Transcranial magnetic stimulation excites the root exit zone of the facial nerve. *Neurosurgery* 1993;32(3):414-6; discussion 415-6.

Toleikis JR, Calvin AO, Shapiro DE, Schafer MF. The use of dermatomal evoked responses during surgical procedures that use intrapedicular fixation of the lumbosacral spine. *Spine* 1993;18(16):2401-7.

Toleikis JR, Skelly JP, Calvin AO, Burkus JK. Spinally elicited peripheral nerve responses are sensory rather than motor. *Clin Neurophysiol* 2000;111(4):736-42.

Toleikis JR, Skelly JP, Calvin AO, Toleikis SC, Bernard TN, Burkus JK, Burr ME, Dorchak JD, Goldman MS, Walsh TR. The usefulness of electrical stimulation for assessing pedicle screw placements. *J Spinal Disord* 2000;13(4):283-9.

Toleikis JR, American Society of Neurophysiological Monitoring. Intraoperative monitoring using somatosensory evoked potentials. A position statement by the American Society of Neurophysiological Monitoring. *J Clin Monit Comput*. 2005;19(3):241-58.

Trammell TR, Friedlander JK, Reed DB. The effect of lower limb ischemia on somatosensory evoked potentials during spinal surgery. Report of two cases and review of the literature. *Spine* 1993;18(3):413-5.

Tsai RY, Yang RS, Nuwer MR, Kanim LE, Delamarter RB, Dawson EG. Intraoperative dermatomal evoked potential monitoring fails to predict outcome from lumbar decompression surgery. *Spine* 1997;22(17):1970-5.

Turkof E, Millesi H, Turkof R, Pfundner P, Mayr N. Intraoperative electroneurodiagnostics (transcranial electrical motor evoked potentials) to evaluate the functional status of anterior spinal roots and spinal nerves during brachial plexus surgery. *Plast Reconstr Surg* 1997;99(6):1632-41.

Ubags LH, Kalkman CJ, Been HD, Drummond JC. The use of a circumferential cathode improves amplitude of intraoperative electrical transcranial myogenic motor evoked responses. *Anesth Analg* 1996;82(5):1011-4.

Ubags LH, Kalkman CJ, Been HD, Porsius M, Drummond JC. The use of ketamine or etomidate to supplement sufentanil/N₂O anesthesia does not disrupt monitoring of myogenic transcranial motor evoked responses. *J Neurosurg Anesthesiol* 1997;9(3):228-33.

Ubags LH, Kalkman CJ, Been HD, Drummond JC. Differential effects of nitrous oxide and propofol on myogenic transcranial motor evoked responses during sufentanil anaesthesia. *Br J Anaesth* 1997;79(5):590-4.

Ubags LH, Kalkman CJ, Been HD. Influence of isoflurane on myogenic motor evoked potentials to single and multiple transcranial stimuli during nitrous oxide/opioid anesthesia. *Neurosurgery* 1998;43(1):90-4; discussion 94-5.

Ubags LH, Kalkman CJ, Been HD, Koelman JH, Ongerboer de Visser BW. A comparison of myogenic motor evoked responses to electrical and magnetic transcranial stimulation during nitrous oxide/opioid anesthesia. *Anesth Analg* 1999;88(3):568-72.

Uematsu S, Tolo V. Electrophysiological recordings during spinal surgery. *Stereotact Funct Neurosurg* 1989;52(2-4):145-56.

Ueta T, Owen JH, Sugioka Y. Effects of compression on physiologic integrity of the spinal cord, on circulation, and clinical status in four different directions of compression: posterior, anterior, circumferential, and lateral. *Spine* 1992;17(8 Suppl):S217-26.

Ulkatan S, Neuwirth M, Bitan F, Minardi C, Kokoszka A, Deletis V. Monitoring of scoliosis surgery with epidurally recorded motor evoked potentials (D wave) revealed false results. *Clin Neurophysiol.* 2006;117(9):2093-101.

van Dongen EP, ter Beek HT, Schepens MA, Morshuis WJ, Langemeijer HJ, Kalkman CJ, Boezeman EH. The influence of nitrous oxide to supplement fentanyl/low-dose propofol anesthesia on transcranial myogenic motor-evoked potentials during thoracic aortic surgery. *J Cardiothorac Vasc Anesth* 1999;13(1):30-4.

van Dongen EP, ter Beek HT, Schepens MA, Morshuis WJ, Langemeijer HJ, de Boer A, Boezeman EH. Within-patient variability of myogenic motor-evoked potentials to multipulse transcranial electrical stimulation during two levels of partial neuromuscular blockade in aortic surgery. *Anesth Analg* 1999;88(1):22-7.

van Dongen EP, ter Beek HT, Schepens MA, Morshuis WJ, de Boer A, Aarts LP, Boezeman EH. Effect of nitrous oxide on myogenic motor potentials evoked by a six pulse train of transcranial electrical stimuli: a possible monitor for aortic surgery. *Br J Anaesth* 1999;82(3):323-8.

van Dongen EP, ter Beek HT, Schepens MA, Morshuis WJ, Haas FJ, de Boer A, Boezeman EH, Aarts LP. The relationship between evoked potentials and measurements of S-100 protein in cerebrospinal fluid during and after thoracoabdominal aortic aneurysm surgery. *J Vasc Surg* 1999;30(2):293-300.

van Dongen EP, ter Beek HT, Aarts LP, Schepens MA, Morshuis WJ, Benning FJ, de Boer A, Boezeman EH. The effect of two low-dose propofol infusions on the relationship between six-pulse transcranial electrical stimulation and the evoked lower extremity muscle response. *Acta Anaesthesiol Scand* 2000;44(7):799-803.

van Dongen EP, Schepens MA, Morshuis WJ, ter Beek HT, Aarts LP, de Boer A, Boezeman EH. Thoracic and thoracoabdominal aortic aneurysm repair: use of evoked potential monitoring in 118 patients. *J Vasc Surg.* 2001;34(6):1035-40.

Vogel LC, Lubicky JP. Neurologic and vascular complications of scoliosis surgery in patients with Ehlers-Danlos syndrome. A case report. *Spine* 1996;21(21):2508-14.

Vossler DG, Stonecipher T, Millen MD. Femoral artery ischemia during spinal scoliosis surgery detected by posterior tibial nerve somatosensory-evoked potential monitoring. *Spine* 2000;25(11):1457-9.

Wada T, Miyamoto T, Murata H, Maeda S, Yamashita K, Yao H. Reconstructive surgery in 59-year-old patient with coarctation of aorta under the monitoring of somatosensory evoked potential and spinal cord perfusion pressure. *Nippon Kyobu Geka Gakkai Zasshi.* 1992;40(1):134-40.

Wada T, Yao H, Miyamoto T, Mukai S, Yamamura M. Prevention and detection of spinal cord injury during thoracic and thoracoabdominal aortic repairs. *Ann Thorac Surg.* 2001;72(1):80-4; discussion 85.

Wass CT, Grady RE, Fessler AJ, Cascino GD, Lozada L, Bechtle PS, Marsh WR, Sharbrough FW, Schroeder DR. The effects of remifentanyl on epileptiform discharges during intraoperative electrocorticography in patients undergoing epilepsy surgery. *Epilepsia.* 2001;42(10):1340-4.

Watanabe K, Watanabe T, Takahashi A, Saito N, Hirato M, Sasaki T. Transcranial electrical stimulation through screw electrodes for intraoperative monitoring of motor evoked potentials. Technical note. *J Neurosurg.* 2004;100(1):155-60.

Watt JW, Fraser MH, Soni BM, Sett PK, Clay R. Total i.v. anaesthesia for transcranial magnetic evoked potential spinal cord monitoring. *Br J Anaesth* 1996;76(6):870-1.

Watt JW, Fraser MH, Soni BM, Sett PK, Clay R. Differential effects of nitrous oxide and propofol on myogenic motor-evoked responses. *Br J Anaesth* 1998;81(2):296.

Weigang E, Hartert M, von Samson P, Pechstein U, Genstorfer J, Pitzer K, Zentner J, Beyersdorf F. Improved spinal cord perfusion during thoracoabdominal aortic repair. *Thorac Cardiovasc Surg.* 2005;53(2):69-73.

Weigang E, Hartert M, Sircar R, V Samson P, Pitzer K, Genstorfer J, Zentner J, Beyersdorf F. Setup of neurophysiological monitoring with tcMEP/SSEP during thoracoabdominal aneurysm repair. *Thorac Cardiovasc Surg.* 2005;53(1):28-32.

Weigang E, Hartert M, von Samson P, Sircar R, Pitzer K, Genstorfer J, Zentner J, Beyersdorf F. Thoracoabdominal Aortic Aneurysm Repair: Interplay of Spinal Cord Protecting Modalities. *Eur J Vasc Endovasc Surg.* 2005; Jul 12 2005; [Epub ahead of print].

Weigang E, Hartert M, Siegenthaler MP, Pitzer-Hartert K, Luehr M, Sircar R, von Samson P, Beyersdorf F. Neurophysiological monitoring during thoracoabdominal aortic endovascular stent graft implantation. *Eur J Cardiothorac Surg.* 2006;29(3):392-6.

Weiss DS. Spinal cord and nerve root monitoring during surgical treatment of lumbar stenosis. *Clin Orthop* 2001;(384):82-100.

Welch WC, Rose RD, Balzer JR, Jacobs GB. Evaluation with evoked and spontaneous electromyography during lumbar instrumentation: a prospective study. *J Neurosurg* 1997;87(3):397-402.

Wiedemayer H, Schaefer H, Armbruster W, Miller M, Stolke D. Observations on intraoperative somatosensory evoked potential (SEP) monitoring in the semi-sitting position. *Clin Neurophysiol.* 2002;113(12):1993-7.

Wiedemayer H, Sandalcioglu IE, Armbruster W, Regel J, Schaefer H, Stolke D. False negative findings in intraoperative SEP monitoring: analysis of 658 consecutive

neurosurgical cases and review of published reports. *J Neurol Neurosurg Psychiatry*. 2004;75(2):280-6.

Williamson JB, Galasko CS. Spinal cord monitoring during operative correction of neuromuscular scoliosis. *J Bone Joint Surg Br* 1992;74(6):870-2.

Wilson-Holden TJ, Padberg AM, Lenke LG, Larson BJ, Bridwell KH, Bassett GS. Efficacy of intraoperative monitoring for pediatric patients with spinal cord pathology undergoing spinal deformity surgery. *Spine* 1999;24(16):1685-92.

Wilson-Holden TJ, Padberg AM, Parkinson JD, Bridwell KH, Lenke LG, Bassett GS. A prospective comparison of neurogenic mixed evoked potential stimulation methods: utility of epidural elicitation during posterior spinal surgery. *Spine* 2000;25(18):2364-71.

Winter RB. Neurologic safety in spinal deformity surgery. *Spine* 1997;22:1527-1533

Witdoeck C, Ghariani S, Guerit JM. Somatosensory evoked potential monitoring in carotid surgery. II. Comparison between qualitative and quantitative scoring systems. *Electroencephalogr Clin Neurophysiol* 1997;104(4):328-32.

Wolfe DE, Drummond JC. Differential effects of isoflurane/nitrous oxide on posterior tibial somatosensory evoked responses of cortical and subcortical origin. *Anesth Analg* 1988;67(9):852-9.

Woodforth IJ, Hicks RG, Crawford MR, Stephen JP, Burke DJ. Electroencephalographic evidence of seizure activity under deep sevoflurane anesthesia in a nonepileptic patient. *Anesthesiology* 1997;87(6):1579-82.

Worth RM, Markand ON, DeRosa GP, Warren CH. Intraoperative somatosensory evoked response monitoring during spinal cord surgery. *Adv Neurol* 1982;32:367-7.

Yamada H, Transfeldt EE, Tamaki T, Torres F, Iazzo PA. The effects of volatile anesthetics on the relative amplitudes and latencies of spinal and muscle potentials evoked by transcranial magnetic stimulation. *Spine* 1994;19(13):1512-7.

Yamada H, Transfeldt EE, Tamaki T, Nishiura H, Taylor BA, Torres F, Iazzo PA. General anesthetic effects on compound muscle action potentials elicited by single or dual spinal cord stimulation. *J Spinal Disord* 1995;8(2):157-62.

Yamamoto N, Takano H, Kitagawa H, Kawaguchi Y, Tsuji H, Uozaki Y. Monitoring for spinal cord ischemia by use of the evoked spinal cord potentials during aortic aneurysm surgery. *J Vasc Surg* 1994;20(5):826-33.

Yamamoto T, Katayama Y, Nagaoka T, Kobayashi K, Fukaya C. Intraoperative monitoring of the corticospinal motor evoked potential (D-wave): clinical index for postoperative motor function and functional recovery. *Neurol Med Chir (Tokyo)*. 2004;44(4):170-80.

Yang LH, Lin SM, Lee WY, Liu CC. Intraoperative transcranial electrical motor evoked potential monitoring during spinal surgery under intravenous ketamine or etomidate anaesthesia. *Acta Neurochir (Wien)* 1994;127(3-4):191-8.

York DH, Chabot RJ, Gaines RW. Response variability of somatosensory evoked potentials during scoliosis surgery. *Spine* 1987;12(9):864-76.

Zentner J, May J. Intraoperative monitoring with somatosensory evoked potentials in neurosurgical operations on the spinal cord. *Cent Nerv Syst Trauma* 1987;4(3):197-207.

Zentner J, Kiss I, Ebner A. Influence of anesthetics--nitrous oxide in particular--on electromyographic response evoked by transcranial electrical stimulation of the cortex. *Neurosurgery* 1989;24(2):253-6.

Zentner J. Noninvasive motor evoked potential monitoring during neurosurgical operations on the spinal cord. *Neurosurgery* 1989;24:709-712

Zentner J. Motor evoked potential monitoring during neurosurgical operations on the spinal cord. *Neurosurg Rev* 1991;14(1):29-36.

Zentner J, Albrecht T, Heuser D. Influence of halothane, enflurane, and isoflurane on motor evoked potentials. *Neurosurgery* 1992;31(2):298-305.

Zentner J, Thees C, Pechstein U, Scheufler KM, Wurker J, Nadstawek J. Influence of nitrous oxide on motor-evoked potentials. *Spine* 1997;22(9):1002-6.

Zhou HH, Turndorf H. Hyper- and hypoventilation affects spinal motor neuron excitability during isoflurane anesthesia. *Anesth Analg.* 1998;87(2):407-10.

Zhou HH, Jin TT, Qin B, Turndorf H. Suppression of spinal cord motoneuron excitability correlates with surgical immobility during isoflurane anesthesia. *Anesthesiology.* 1998;88(4):955-61.

Zhou HH, Zhu C. Comparison of isoflurane effects on motor evoked potential and F wave. *Anesthesiology* 2000;93(1):32-8.

Zhou HH, Kelly PJ. Transcranial electrical motor evoked potential monitoring for brain tumor resection. *Neurosurgery.* 2001;48(5):1075-80.

Zileli M, Taniguchi M, Cedzich C, Schramm J. Vestibulospinal evoked potential versus motor evoked potential monitoring in experimental spinal cord injuries of cats. *Acta Neurochir (Wien)* 1989;101(3-4):141-8.

Zornow MH, Drummond JC. Intraoperative somatosensory evoked responses recorded during onset of the anterior spinal artery syndrome. *J Clin Monit.* 1989;5(4):243-5.